

## Herbal, Animal and Mineral Remedies in Burn Wound: a review of Persian Traditional Medicine literature

Mahboobeh Raeiszadeh <sup>1,2</sup>, Nasser Ebrahimpour <sup>3,4</sup>, Maryam Iranpour <sup>5</sup>, Mitra Mehrabani <sup>6</sup>, Mehrnaz Mehrabani <sup>7</sup>, Zeinab Kordestani <sup>8</sup>, Mehrzad Mehrbani <sup>6\*</sup>

1. Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran
2. Department of Traditional Pharmacy, Faculty of Traditional Medicine, Kerman University of Medical Sciences, Kerman, Iran
3. Pharmaceutical Sciences and Cosmetic Products Research Center, Kerman University of Medical Sciences, Kerman, Iran
4. Department of Traditional Medicine, Faculty of Traditional Medicine, Kerman University of Medical Sciences, Kerman, Iran
5. Pathology and Stem Cell Research Center, Kerman University of Medical Sciences, Kerman, Iran
6. Herbal and Traditional Medicines Research Center, Kerman University of Medical Sciences, Kerman, Iran
7. Physiology Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran
8. Cardiovascular Research Center, Institute of Basic and Clinical Physiology Sciences, Kerman University of Medical Sciences, Kerman, Iran



### ABSTRACT

**Background:** Burn wound is one of the most common injuries worldwide. However, its management still remains a health concern and research is ongoing for more efficient therapies. Persian medicine has described different methods in this regard. Previous reviews have mostly focused on the herbal-based therapies for burn management. Hence, the current study aimed to review both non-herbal and herbal-based therapies used in Persian medicine for their burn healing properties, and to compare them with new evidence.

**Method:** The most important texts of Persian medicine were reviewed for burn wound classification, managements and treatment approaches. Moreover, herbal, minerals, and animal-based products claimed to possess burn wound healing activity, were extracted. In the next step, digital databases were searched to find new supporting data for the extracted-traditional remedies and their healing mechanisms.

**Results:** More than one hundred traditional medicinal herbs, minerals, and animal-based products have been recommended to treat burn wound in Persian medicine among them, the healing effect of thirty-five herbs and eight non-herbal treatments (mineral and animal-based compounds) were supported by new evidence. Otherwise, there was not new evidence reporting the healing effect of seventy-three herbs and sixteen non-herbal compounds extracted from Persian Medicine literature.

**Conclusion:** Although some managements of burn wounds in Persian medicine have been confirmed by new evidence, the efficacy of a plenty of materials needs to be methodically evaluated. Hence, it could introduce new clues for future research.

**Keywords:** Burn wound, Persian medicine, Medicinal herbs, Minerals, Animal-based products

**Citation:** Raeiszadeh M, Ebrahimpour N, Iranpour M, Mehrabani M, Mehrabani M, Kordestani Z, Mehrbani M. Herbal, Animal and Mineral Remedies in Burn Wound: a review of Persian Traditional Medicine literature. *Journal of Kerman University of Medical Sciences* 2021; 28(4): 520-538. doi: 10.22062/JKMU.2021.91771

**Received:** 19.08. 2020

**Accepted:** 11.10. 2021

**\*Correspondence:** Mehrzad Mehrbani; Email: m.mehrbani@kmu.ac.ir

Published by Kerman University of Medical Sciences

## Introduction

**B**urning is the most devastating injury with a high potential for morbidity and mortality (1). It was estimated that, 6 million patients per year need medication for burns worldwide, but most of them are treated in outpatient clinics (2). However, burn shock, multi-organ failure, local and systemic resistant infections, hypertrophic scars and keloids, chronic wounds, cosmetic attentions, underlying disease and special conditions like pregnancy and childhood are serious challenges (3-6).

Rapid wound healing is considered the main aim of burn management to minimize infection and decrease functional and aesthetic complications. Essentially, topical medication has been used to diminish morbidity and mortality in patients with severe burns (7).

There are several topical medications in the market for the management of burn wounds. However, some of the most common drugs could have antimicrobial activity rather than wound healing effect with considerable side effects (8, 9). Although using tissue scaffolds, nanoparticles, reproductive stimulatory factors, stem cells, and gene therapy are relatively modern treatments, but more studies should be conducted for cost-effective methods (10).

Over the centuries, people have had to use the materials around them as medications and gradually, they have partially refined more effective subjects (11-17). The texts of traditional Persian medicine (TPM) possess a high investigational potential in introducing new methods and medications (18, 19). TPM literature has described some procedures for burn management and also introduced many compounds of plant, mineral and animal origin.

The current study has reviewed the perspective of TPM on burn wounds managing and introduces numerous herbal, animal, and mineral materials used from the 10<sup>th</sup> to 19<sup>th</sup> centuries, with their effects on wound healing that have been reported in the recent studies.

## Material and Method

The most important texts of TPM, including the Canon of Medicine (20), Zakhire Kharazmshahi (21), Al-Hawi (22), Al-Mansouri

fi al-Tibb (23) and Makhzan al-Advieh (24), were reviewed for burn wound classification, managements and treatment approaches. Moreover, medicinal herbs, minerals, and animal-based products, claimed to have burn wound healing activity, were extracted. Then, electronic databases, including Google Scholar, PubMed, and Scopus were searched for the related evidence from 2010 to 2019, using the search terms of “burn” or “wound”, and the “scientific name” or “common name” of the materials in the title or abstract. All the collected articles were evaluated for inclusion of any *in-vivo*, animal, or clinical evidence of their efficacy related to burn and wound healing properties. Only articles in English language were included in the review. Review articles, case reports, and *in-vitro* studies were excluded. If a subject was not found in any study in the current time range, the search strategy was expanded to the previous years. If there had been many studies on the same topic, such as *Calendula officinalis*, some of them would have presented (researchers' discretion). The included articles were reviewed based on the used part of the component (only herbs), having single or combined formula, type of wound, and the type of the investigational model (i.e. animal or human). Also, the evidence was reviewed for histological (histopathology and immunohistochemistry tissue markers), molecular (oxidative and antioxidative markers and gene expressions), macroscopic (wound closure rate), mechanical (stretch test), physical (odor, scab, bed color and, exudate), and biological (antimicrobial or antifungal) parameters.

## Results

### 1.1. Definitions and classification of wounds in TPM

In TPM, wound is defined as discontinuity of the skin, caused by external damage (hit, cut, burn, etc.), lesions under the skin (abscess, tumors, etc.), and skin rashes. Also, to determine the greatest treatment, wounds are described according to the following characteristics (Table 1).

**Table 1.** Defining Wound characteristics (20, 21, 25)

Characteristic	Classification
Cause	Hit/ burn/ abscess/ infection/ malignancy
Extent	Wide/ small
Location	Skin or mucus / involved area like face or organs
Depth (severity)	With or without blisters, along with damage to muscles or bones
Exudate	Color, odor, thickness, amount
Underlying health condition	Presence of underlying or associated illness, patient's tolerance against disease, age
Surrounding area	Swelling, torsion, inflammation, pain
Margins	Thickening, redness, inflammation, pain
Extension (conversion)	Infection, malignity
Weather considerations	Temperature, humidity, intensity and direction of wind

In TPM, burn is categorized as a separate group of injuries, and it is classified according to causative agents of fire, hot appliance, hot water, hot oil, chemicals, sunshine, and thunder.

### 1.2. Burned patient management in TPM

1. Early management of burn included cooling and cleaning the wound using cold

water. Topical treatments were used to reduce pain and prevent blisters (Table 2) (20).

Lime salve, which is a combination of calcium hydroxide powder with herbal oils such as sesame or olive oils, is a common topical treatment used for burn wounds (26).

**Table 2.** Topical medications used to reduce pain and prevent blisters in TPM

Components
Red powdered Sandal wood ( <i>Pterocarpus santalinus</i> L.), areal part of Betel nut ( <i>Areca catechu</i> L.) and earthenware
The extract of Black nightshade ( <i>Solanum nigrum</i> ), rose flower ( <i>Rosa Damascena</i> ) distillate and powdered earthenware
Oily extract of rose flower and egg yolk
Oily extract of rose flower, egg yolk, washed barely flour and extract of aerial part of chicory ( <i>Cichorium intybus</i> )
Baked lentils ( <i>Lens culinaris</i> Medik.) and oily extract of rose flower
Armani <sup>®</sup> soil (a soil containing some heavy metals like Iron, Copper and Lead) and vinegar
Washed Lime (Calcium Hydroxide), egg white, Sefidab (a white-colored substance containing tin, silver or zinc components), opium, oily extract of rose flower, beeswax and milk
The extract of Malva ( <i>Malva sylvestris</i> ), black nightshade ( <i>Solanum nigrum</i> ) and coriander ( <i>Coriandrum sativum</i> L.), Sefidab, Mordarseng (a soil containing heavy metals like Iron, Copper, and Lead) and the oily extract of rose flower

2. After burn injury, especially a large or severe one, a diet including light foods and nourishing fluids was being used (21-23, 26, 27):

- Oxymel (a syrup of vinegar and honey)
- Herbal extracts of Mallow (*Malva sylvestris*), Chicory (*Cichorium intybus*), and water lily (*Nymphaea odorata*)
- Juices of jujube (*Ziziphus jujuba* Mill.), cherry (*Prunus avium*), purslane (*Portulaca oleracea*), plum (*Prunus domestica*), and pomegranate (*Punica granatum*)
- Easy-to-digest foods; barley soup containing vegetables and little bird meat like partridge, soft-boiled eggs and sugar-almond syrup.
- After a few days and with the onset of congestion, it is recommended to eat heavier and nutritious foods, such as broth made with meats, beans and spices.

3. To relieve pain, two treatments were being used (20, 23):

- Topical supplements containing vegetable oils and egg white

- Oral drugs such as opium (*Opium poppy*), Mandrake (*Mandragora officinarum*) flower and poppy (*Papaver somniferous*) seeds

4. Other procedures of TPM in the management of burn wounds were Fasd (phlebotomy) and Hijamat (cupping) (28). Fasd is a special practice in TPM performed by fine cutting of a superficial vein and extracting some blood for therapeutic purposes (20). Usually, blood is taken from the veins away from inflamed site to control pain and inflammation; for example, phlebotomy of the left basilic vein in the right upper limb injury (29). Hijamat includes taking blood from the surface of the skin by cupping. TPM physicians believe that this practice helps to control inflammation (20, 29).

5. Surgery operations (21, 25):

- Excessive scarring (hypertrophic scar and keloid)

- Cutting the bed of wound to create a way out of the secretions

- Debridement, refreshment of the wound edges, and the removal of necrotic bones and other necrotic tissues

6. Compression bandage, wound drainage with oily wicks of cotton

7. Treating fever and other general complications

8. Treatment and prevention of injury-disability and rehabilitation with local softening agents, such as almond or olive oil with spices, massage and surgery (20, 22)

1.3. Materials used in burn wounds in TPM and new related evidences

Based on the TPM references mentioned in the method section, more than one hundred traditional medicinal herbs, minerals and animal-based products have been recommended to treat a burn wound. As shown in Table 3, most of the herbs that new evidences show their effects in burn wound healing are from Asteraceae, Boraginaceae, and Leguminosae family. Moreover, most of the results were related to fenugreek, marigold, arnebia, St

John's wort, and sesame, in the order of their appearance. According to the 84 articles found on plants used for burn wound healing, the most used parts of herbs were leaf (26 cases), seed (21 cases), and flower (19 cases). Also, the other used parts of herbs were stem, root, bark, aerial part, fruit as well as resin. In the herbal studies on burn and wound healing, most of the *in-vivo* models were excision, incision, and burn wound models. Furthermore, among the 84 articles reviewed, the animals used in burn and wound healing models were rat (60 cases), mice (11 cases), human (6 cases), rabbit (4 cases), pig (2 cases), and dog (1 case). In non-human studies, the most studied parameters were included histological and macroscopic evaluations. Also, in human studies, macroscopic and physical parameters were often assessed.

As demonstrated in tables 4 and 5, fewer studies have been conducted on mineral and animal-based materials and most studies included silver, zinc, and honey. Despite the application of some materials in topical burn management in TPM, there was no new study in the databases (Tables 6 and 7). Therefore, new studies can be conducted on these materials such as medicinal herbs belonged to 47 families, particularly Asteraceae and Fabaceae families.

**Table 3.** Some new evidence of the topical medicinal herbs used in TPM for burn wound healing

Family name	Scientific name	Common name	Traditional name	Part used	Single/combination	Study model	Animal type	Study parameters				
								Histological	Molecular	Macroscopic	Mechanical	Physical
Asphodelaceae	<i>Aloe vera</i> L.	Aloe vera	Sebr	Leaf	S	Chronic ulcer (30)	Human			*		*
				Leaf	S	Burn wound (31)	Human					*
				Leaf	S	Burn wound (32)	Rat	*		*		
Asparagaceae	<i>Dracaena cochinchinensis</i> Lour.	Dragon's blood	Dam-ol-akhavain	Resin	S	Excision/incision wound (33)	Rat			*	*	
				<i>Calendula officinalis</i> L.	Marigold	Hai-ol-aalam	Flower	S	Excision wound (34)	Rat	*	*
Asteraceae				Flower	C	Excision/ burn wound (35)	Mice			*		*
				Flower	S	Excision wound (36)	Rat	*	*	*	*	*
				Flower	C	Excision wound (37)	Rabbit	*	*	*	*	*
				Flower	S	Excision wound (38)	Rat	*		*		
				Flower	S	Diabetic wound (39)	Human				*	*
	<i>Cichorium intybus</i> L.	Chicory	Hendebe	All organs	S	Incision/ excision wound (41)	Rat	*		*		

**Table 3.** Some new evidence of the topical medicinal herbs used in TPM for burn wound healing

	<i>Tragopogon graminifolius</i> DC.		Sheng	leaf and stem	S	Burn wound (42)	Rat	*	*	*
				Leaf & root	S	Burn wound (43)	Rat	*	*	
				Leaf	S	Burn wound (44)	Rat	*	*	* *
Boraginaceae	<i>Arnebia euchroma</i> L.	Arnebia	Obukhalsa	Leaf	S	Burn wound (43)	Rat	*	*	
				Leaf	S	Excision wound (45)	Rat	*	*	
				Bark resin	S	Burn wound (46)	Rat		*	*
				Leaf	C	Burn wound (47)	Rat	*	*	
				Flower	C	Excision wound (48)	Rat	*	*	
Brassicaceae	<i>Brassica oleracea</i> L.	Turnip	Shaljam	Flower	S	Excision wound (49)	Rat	*		
				Flower	C	Excision wound (50)	Rat	*	*	
Burseraceae	<i>Boswellia carteri</i> L.	Frankincense (olibanum)	Kondor	Resin	C	Excision wound (51)	Mice	*	*	*
Fagaceae	<i>Quercus sp.</i>	Oak	Afes	Flower	S	Burn wound (52)	Pig		*	*
				Flower	S	Excision wound (53)	Dog	*	*	*
				Aerial parts	S	Incision wound (54)	Rat	*	*	*
Hypericaceae	<i>Hypericum perforatum</i> L.	St Johns wort	Hofarighon	All organs	S	Excision wound (55)	Rat	*	*	
				Flower	S	Burn wound (56)	Rat	*		
				Aerial parts	S	Excision/ incision/ burn wound (57)	Rat	*	*	
				Leaf	S	Excision wound (58)	Rat	*	*	*
Iridaceae	<i>Crocus sativus</i> L.	Saffron	Zafaran	Flower	S	Burn wound (59)	Mice	*	*	
	<i>Ocimum basilicum</i> L.	Sweet basil	Shahasfarm bostani	Leaf	C	Excision wound (60)	Rat		*	
				Leaf	S	Incision/ excision wound (61)	Rat	*	*	* *
Lamiaceae	<i>Ocimum sanctum</i> L.	Holy basil	Shahasfarm jabali	Leaf	S	Excision wound (62)	Rat	*	*	
				Leaf	S	Incision/excision wound (63)	Rabbit		*	*
Lauraceae	<i>Cinnamomum comphora</i> L.	Camphor	Kafur	Bark resin	C	Burn wound (64)	Rat	*	*	
	<i>Acacia auriculiformis</i> A.Cunn.	Ear leaf acacia	Aghagheia	Stem bark	S	Excision /incision wound (65)	Mice	*	*	* * *
	<i>Acacia leucophloea</i> Roxb.	White bark acacia	Aghagheia	Bark	S	Excision/ incision wound (66)	Rat	*	*	* * *
	<i>Ceratonina siliqua</i> L.	Locust bean (carob)	Kharnub	Bean	C	Excision wound (67)	Pig	*	*	
Leguminosae				Seed	S	Excision/ incision/ dead space wound (68)	Rat		*	* *
				Seed	S	Excision wound (69)	Rat		*	* *
	<i>Trigonella foenum graecum</i> L.	Fenugreek	Holbeh	Seed	C	Excision wound (70)	Rat		*	*
				Seed	S	Excision wound (71)	Mice	*	*	
				Seed	C	Excision wound (72)	Rat	*	*	* * *

**Table 3.** Some new evidence of the topical medicinal herbs used in TPM for burn wound healing

				Seed	C	Excision wound (73)	Rat	*	*	*
				Seed	S	Incision wound (74)	Rabbit	*		
				Seed	S	Incision/excision/dead space wound (75)	Rat/Mice	*	*	*
				Seed	C	Excision wound (76)	Rat	*	*	*
				Seed	S	Excision wound (77)	Rat	*		
Linaceae	<i>Linum usitatissimum</i> L.	Flax seed	Katan	Seed	S	Excision/ incision wound (78)	Rat	*	*	*
				Seed	C	Excision/ incision wound (79)	Rat		*	*
				Leaf	C	Excision wound (80)	Rat	*	*	*
Lythraceae	<i>Lawsonia inermis</i> L.	Henna	Hana	Leaf	S	Incision wound (81)	Rat		*	
				Leaf	S	Excision/incision/ dead space wound (82)	Rat	*	*	*
				Flower	S	Incision wound (83)	Mice	*		
Malvaceae	<i>Malva sylvestris</i> L.	Malva	Khobbazi	Flower	C	Excision wound (84)	Rat	*	*	
				Flower	C	Burn wound (85)	Rat	*	*	*
				Leaf	C	Excision/incision wound (86)	Rat	*	*	*
Meliaceae	<i>Azadirachta indica</i> A.Juss.	Neem	Azad derakht	Seed	S	Non-healing chronic wound (87)	Human	*	*	*
				Stem bark	S	Excision wound (88)	Mice		*	*
Moraceae	<i>Morus alba</i> L.	Mulberry	Toot	Leaf	C	Excision/ dead space wound (89)	Rat	*	*	*
Musaceae	<i>Musa paradisiaca</i> L.	Banana	Moze	Leaf	S	Burn wound (90)	Rat	*	*	*
Myrtaceae	<i>Myrtus communis</i> L.	Myrtle	Aas	Leaf	S	Burn wound (91)	Rat		*	*
				Leaf, stem and seed	S	Incision wound (92, 93)	Rat	*	*	
				Fruit	S	Burn wound (94)	Human		*	*
Oleaceae	<i>Olea europaea</i> L.	Olive	Zaitoon	Leaf	S	Incision wound (95)	Rat	*		*
				Fruit	C	Burn wound (96)	Mice	*	*	*
				Fruit	C	Burn wound (97)	Mice	*	*	*
				Barry and cone	S	Incision /excision wound (98)	Rat	*	*	*
				Seed	S	Excision / incision / dead space burn wound (99)	Rat	*	*	*
				Seed	S	Excision / incision/ dead space wound (100)	Rat		*	*
Pedaliaceae	<i>Sesamum indicum</i> L.	Sesame	Semsem	Seed	S	Incision wound (101)	Rat		*	
				Seed	S	Burn wound (102)	Rat	*	*	
				Seed	C	Burn wound (103)	Rat	*	*	*
Plantaginaceae	<i>Plantago major</i> L.	Broadleaf plantain	Bazr-e-ghatona	Seed	S	Burn wound (104)	Rat	*		*
				Leaf	S	Cutaneous incision wound (105)	Mice	*	*	*

**Table 3.** Some new evidence of the topical medicinal herbs used in TPM for burn wound healing

Family name	Scientific name	Traditional name	Part	Preparation	Study model	Animal type	Histological	Biochemical	Macroscopic	Mechanical	Physical	Biological
Rosaceae	<i>Plantago ovate</i> L.	Psyllium	Bazr-eghatona	Seed husk	S	Excision wound (106)	Rat	*	*	*	*	*
	<i>Cydonia oblonga</i> L.	Quince seed	Safarjal	Seed	S	Toxin-induced wound (107)	Rabbit	*				*
				Seed	S	Biopsy punch (108)	Human			*		
	<i>Rosa Damascena</i> L.	Damask rose	Vard	Flower	C	Excision wound (109)	Rat	*		*		
Flower				C	Burn wound (85)	Rat	*	*				*
Rutaceae	<i>Citrus limon</i> L.	Lemon	Limou	Fruit	S	Incision wound (110)	Mice	*		*		
Solanaceae	<i>Solanum nigrum</i> L.	Nightshade	Enab-ol-salab	Leaf	C	Burn wound (85)	Rat	*	*			*
Tamaricaceae	<i>Tamarix aphylla</i> L.	Saltcedar	Tarfa	Leaf	S	Excision wound (111)	Rat			*		

**Table 4.** Some new evidence of the topical minerals used in TPM for burn wound healing

Name	Molecular formula	Single/combination	Study model	Animal type	Study parameters					
					Histological	Biochemical	Macroscopic	Mechanical	Physical	Biological
Silver	Ag	S	Burn wound (112)	Human			*		*	
		S	Pressure ulcers (113)	Human			*		*	
		S	Burn wound (114)	Rat					*	*
		S	Burn wound (115)	Human					*	*
		S	Donor site (116)	Human					*	*
Zinc	Zn	S	Burn wound (117)	Rat		*	*			
		S	Burn wound (118)	Rat	*		*			*
		S	Burn wound (119)	Rabbit	*		*		*	
Lime	Ca(OH) <sub>2</sub>	S	Excision wound (120)	Human			*		*	
		C	Burn wound (96)	Mice	*	*				*
Copper	Cu	C	Burn wound (97)	Rat					*	*
		S	Diabetic wound (121)	Human			*		*	*
		S	Excision wound (122)	Mice	*	*	*			

**Table 5.** Some new evidence of the topical animal-based used in TPM for burn wound healing

Name	Single/combination	Study model	Animal type	Study parameters					
				Histological	Biochemical	Macroscopic	Mechanical	Physical	Biological
Honey	S	Burn wound (123)	Human			*			
		Episiotomy wound (124)	Human			*			
		Incision wound (125)	Human			*			
		Burn wound (126)	Rat	*		*			
		Burn wound (127)	Human			*		*	*
		Cutaneous wound (128)	Rat	*		*			
		Incision/excision wound (129)	Rat	*		*		*	
		Intraoral wound (130)	Rat	*		*		*	
		Excision wound (131)	Mice	*		*		*	
		Incision/ excision/ burn wound (132)	Rat	*		*		*	*
		Burn wound (133)	Human			*		*	*
		Incision wound (134)	Mice	*		*	*	*	*
		Incision wound (135)	Rabbit			*	*	*	*
Spider web	S	Excision/ incision wound (136)	Rat	*		*	*		

**Table 6** Medicinal herbs used in the treatment of burn wounds in TPM, without new evidence

Family name	Traditional name	Scientific name	Common name	Used part
Amaranthaceae	Selgh	<i>Beta vulgaris</i> L.	Beet	Leaf/ root

**Table 6** Medicinal herbs used in the treatment of burn wounds in TPM, without new evidence

Amaryllidaceae	Koras	<i>Allium schoenoprasum subsp. iranicum</i>	Chives	Leaf
Anacardiaceae	Somagh	<i>Rhus coriaria</i> L.	Sumac	Fruit
Apiaceae	Kozborah	<i>Coriandrum sativum</i> L.	Coriander	Leaf/ seed
	Javsheir	<i>Opopanax chironium</i> L.	Sweet myrrh	Resin
Araliaceae	Lablab	<i>Hedera helix</i> L.	Common ivy	Seed
Areceaceae	Fufal	<i>Areca catechu</i> L.	Betel nut	Aerial part
Aristolochiaceae	Zaravand	<i>Aristolochia rotunda</i> L.	Smearwort	Root
Asteraceae	Eshkheis	<i>Atractylis gummifera</i> L.	Stemless atractylis	Root
	Harshaf	<i>Cynara scolymus</i> L.	Artichokes	Root
	Lahiat-ol-tis	<i>Tragopogon pratensis</i> L.	Meadow salsify	Leaf/ root
	Argheitoun	<i>Arctium tomentosum</i> Mill.	Woolly burdock	Root/ seed
	Oghhovan	<i>Tanacetum parthenium</i> L.	Feverfew	Aerial part
Boraginaceae	Kornob	<i>Brassica oleracea</i> L.	Cabbage	Aerial part
	Fojl	<i>Raphanus sativus</i> L.	Radish	Leaf/ seed
Bursaceae	Morr	<i>Commiphora myrrha</i> Engl.	Myrrh	Resin
	Moghl	<i>Commiphora mukul</i> Engl.	Gugal	Resin
Capparaceae	Kabar	<i>Capparis spinosa</i> L.	Caper bush	Root
Cistaceae	Ladan	<i>Cistus ladanifer</i> L.	Labdanum	Resin
Colchicaceae	Surenjan	<i>Colchicum</i> sp.	Autumn crocus	Aerial part
Convolvulaceae	Habb-ol-nil	<i>Ipomoea hederaceae</i> Jacq.	Ivyleaf morningglory	Seed
	Torbod	<i>Silva manso</i> L.	Turpeth	Root
Cucurbitaceae	Ghesa	<i>Cucumis sativus</i> L.	Cucumber	Fruit
	Ghar	<i>Cucurbita pepo</i> L.	Squash	Fruit
Cupressaceae	Abhol	<i>Juniperus sabina</i> L.	Savin juniper	Fruit/ leaf
Dipsacaceae	Mamisa	<i>Scabiosa arvensis</i> L.	Scabiosa	Seed
Ebenaceae	Aabnus	<i>Diospyros ebenum</i> J. Koenig ex Retz.	Ebony	Bark
Fabaceae	Bagham	<i>Caesalpinia sappan</i> L.	Sappanwood	Leaf
	Adas	<i>Lens culinaris</i> Medik.	Lentil	Seed
	Nil	<i>Indigofera tinctoria</i> L.	True indigo	Aerial part
	Baghella	<i>Vicia faba</i> L.	Faba bean	Seed
	Kasira	<i>Astragalus gummifer</i> Labill.	Milkvetch	Leaf
	Sandal	<i>Pterocarpus santalinus</i> L.	Red sandalwood	Wood
	Termes	<i>Lupinus termis</i> Forssk.	White lupin	Seed
	Darshishaan	<i>Calicotome spinosa</i> L.	Spiny broom	Fruit
	Sana makki	<i>Senna acutifolia</i> L.	Alexandrian senna	Leaf
	Karasne	<i>Vicia ervilia</i> L.	Ervil	Seed
Gentianaceae	Jentiana	<i>Gentiana lutea</i> L.	Yellow gentian	Leaf/ root
Lamiaceae	Komazarius	<i>Teucrium chamaedrys</i> L.	Wall germander	Aerial part
Liliaceae	Susan	<i>Lilium candidum</i> L.	Madonna lily	Flower
Loranthaceae	Debgh	<i>Loranthus europaeus</i> Jacq.	Mistletoe	Seed
Malvaceae	Khatmi	<i>Alcea</i> sp.	Hollyhocks	Leaf
	Ghoton	<i>Gossypium herbaceum</i> L.	Levant cotton	Wool
Oxalidaceae	Hammaz	<i>Oxalis acetosella</i> L.	Wood sorrel	Leaf/ root
Palmaceae	Tamr	<i>Phoenix dactylifera</i> L.	Date	Fruit
	Narjil	<i>Cocos nucifera</i> L.	Coconut	Fruit
Papaveraceae	Khashkhash	<i>Papaver somniferum</i> L.	Opium poppy	Seed
Pinaceae	Sanoubar	<i>Pinus albicaulis</i> Engelm	Spruce	Leaf/ root



**Table 6** Medicinal herbs used in the treatment of burn wounds in TPM, without new evidence

Platanaceae	Dolb	<i>Platanus orientalis</i> L.	Oriental plane	Bark/ fruit
Poaceae	Jow	<i>Hordeum vulgare</i> L.	Barley	Seed
	Hentah	<i>Triticum aestivum</i> L.	Wheat	Seed
	Oroz	<i>Oryza sativa</i> L.	Rice	Seed
Polygonaceae	Asioraie	<i>Polygonum aviculare</i> L.	Pigweed	Leaf
Portulacaceae	Baghlah-ol-hamgha	<i>Portulaca oleracea</i> L.	Pursley	Seed
Potamogetonaceae	Jaronahr	<i>Potamogeton natans</i> L.	Broad-leaved pondweed	Leaf
Primulaceae	Anaghalls	<i>Anagallis arvensis</i> L.	Pimpernel	Leaf
Pteridaceae	Barsiavashan	<i>Adiantum capillus-veneris</i> L.	Maidenhair	Aerial part
Punicaceae	Romman	<i>Punica granatum</i> L.	Pomegranate	Fruit
Ranunculaceae	Piaranga	<i>Thalictrum foliolism</i> DC.	Meadow-rue	Root
	Beesh	<i>Aconitum ferox</i> Wall. Ex Ser	Indian Aconite	Root
Rhamnaceae	Annab	<i>Ziziphus jujuba</i> Mill.	Jujube	Leaf
Rosaceae	Loz-ol-morre	<i>Prunus amygdalus var. amara</i>	Bitter almond	Seed
Rubiaceae	Ghalion	<i>Galium aparine</i> L.	Cleavers	Flower
Salicaceae	Gharab	<i>Salix babylonica</i> L.	Babylon willow	Leaf/ flower
Smilacaceae	Choob-e-chini	<i>Smilax china</i> L.	China root	Leaf/ root
	Oshbe maghrebie	<i>Smilax sarsaparilla</i> L.	Honduran	Leaf/ flower
Solanaceae	Osaj	<i>Lycium afrum</i> L.	Honey thorn	Root
	Kakanj	<i>Physalis alkekengi</i> L.	Chinese lantern	Fruit
Violaceae	Banafsaj	<i>Viola tricolor</i> L.	Pansy	Root
Vitaceae	Serkeh	<i>Vitis vinifera</i> L.	Vinegar	Fruit extract
Xanthorrhoeaceae	Ashraas	<i>Eremurus spectabilis</i> M.Bieb.	Eremurus	Root
Zingiberaceae	Jadvar	<i>Curcuma zedoaria</i> Rosc.	Zedoary	Root

**Table 7.** None-herbal materials used in the treatment of burn wounds in TPM, without new evidence

Traditional name	Formulation	Dosage forms
Salt	NaCl	Powder
Sangarf/ osranj/ mordasang	PbO	Powder
Wood ash	-	Powder
Shadanj	Fe <sub>2</sub> O <sub>3</sub>	Powder
Green zadj	FeSO <sub>4</sub>	Powder
Lead cutting	Pb	Powder
Iron cutting	Fe	Powder
Earthenware	-	Powder
Tin whiter	SnO	Powder
Shell	-	Powder
Orpiment	As <sub>2</sub> S <sub>3</sub>	Powder
Spider's web	-	Dried powder
Animal feces	-	Dried powder/ ash
Animal skin	-	Dried powder/ ash/fresh skin as dressing
Animal hair	-	Dried powder/ ash
Elephant tusk	-	Dried powder

## Discussion

Human beings have been familiar with the burn, wound injuries, and their management. Apart from the basics and methods of dealing with burn patients, the main difference between ancient and new therapies is the use of natural substances available in the living environment for treatments. These materials have usually been used in combination and gradually improved in quality and effectiveness. New evidence for traditional remedies is growing

rapidly (137). Many of the herbal, mineral, and animal-based materials used for burn wound healing in TPM have been evaluated in recent years. Our results showed that the most common type of study for evaluating burn-healing materials were animal models, especially rats. Due to the availability, reasonable cost, easier methods, and structural similarities of its skin with human skin, rats are the most suitable animals for burn and wound healing examinations (17, 138). Human models are not

used for initial observations due to the large differences in the types of burns, ethical issues, and the difficulty of matching the samples. On the other hand, *in vitro* studies are limited in providing practical results to provide the basics of clinical studies. Therefore, it seems that using animal models is the best choice in most cases.

Our results showed that most of the herbs that have new evidence in burn wound healing were from the Asteraceae family. The efficacy of several plants from the Asteraceae family as wound healing agents has been indicated in ethnopharmacological studies. Due to distribution and ethnopharmacological importance of the plants, several plant products derived from this family have been studied, which some of their pharmacological activities have been already identified. These pharmacological activities include anti-inflammatory, antimicrobial, antioxidant, anti-protozoa, and healing activities. Also, their wound healing efficacy has been suggested to be due to their ability to promote the proliferation of keratinocytes and thus, the remodeling of the extracellular matrix (139). The Asteraceae species have been identified to produce numerous secondary metabolites, such as polyphenols, flavonoids, diterpenoids, and sesquiterpene lactones (140). *Cynara scolymus* L. and *Tanacetum parthenium* L. are from Asteraceae family used as components of topical remedy for burn wound in TPM. *Cynara scolymus* L. contains considerable amounts of minerals and polyphenolic compounds, along with antioxidant and cardioprotective effects (141-143). Also, numerous studies have argued the pharmacologic properties such as anti-inflammatory effect of *Tanacetum parthenium* L. (144). One member of this family, *Atractylis gummifera* L., has been used as a topical medication for burn healing in TPM. However, since the hepatotoxicity and nephrotoxicity effects of this drug has been reported in previous studies, its frequent topical administration, might lead to liver and renal injury (145).

The herbs of Boraginaceae family are well-known for their pharmacological properties. *Arnebia* species are rich in naphthoquinones, such as alkannins, shikonins, and their derivatives have different biological activities, including anti-inflammatory, antimicrobial, wound-healing, and anti-tumorous effects (43, 146). Borage seeds oil is rich in gamma-linoleic acid, a valuable component in wound healing

owing to its anti-inflammatory effects (147, 148).

An oily amorphous resin obtained from the Burseraceae family chemically contains a mixture of triterpenes, sitostenonein, and mono and sesquiterpenes. Some of these compounds also present anti-inflammatory and antimicrobial activity (149). In TPM, the resins are widely used in topical treatments (150).

Isoflavonoids are the phytochemical characteristics of the Fabaceae, and have a limited distribution in non-leguminous plant families (151). Although there is limited evidence for the therapeutic use of Fabaceae family plants in modern medicine sources, they have been widely used in topical burn compounds in TPM, providing an appropriate research area.

The stool of animals, especially pigeons, has been reported in TPM as a material used for local therapy of burn wounds (21). The noteworthy point in this regard is whether the fecal microbial flora can affect wound healing processes. Several studies have investigated the effects of probiotics on skin disorders and their positive effects have been indicated (152, 153). In a clinical study, bacteria *Lactobacillus plantarum* had the same effect as silver sulfadiazine on the control of bacterial infection of burn wounds (154). In an *in-vitro* study, it was reported that this bacterium could inhibit the proliferation of *Pseudomonas aeruginosa* (155). In another study, *lactobacillus acidophilus* inhibited the bacteria isolated from burn wounds (156). However, there was no study on the effect of stool bacteria on wound pathogens. Other materials mentioned in TPM for the treatment of burn wounds include the skin of animals, such as sheep, goat, and horse, which are transient grafts (covering the surface of the burn with freshly squeezed animal skin), and ash or powdered skin (23). The use of pig xenografts in burn wounds has a successful history (157, 158). Animal collagen has been studied in the treatment of wound and currently is used to make new medications (159, 160). However, due to some issues such as health, environmental and ethical concerns, it is not possible to widely use animal products to produce medicine as has been common in ancient times. But, with the aid of new research, they can be promoted to be used for developing of tissue scaffolds, hormone therapy, and isolation of microbial flora.

Also, the use of rocks and soils containing heavy metals (copper, lead, tin, silver, zinc and

iron) in the combination of topical drugs was common in TPM (22, 23, 161). Among these compounds, silver still has had prominent position in wound treatment (162, 163). Zinc compounds are also used to treat various types of wounds (164, 165). However, in animal and human studies, copper-based coatings are effective in the treatment of wounds (121, 122). The use of tin and lead compounds has been obsolete, due to the side effects and systemic absorption. However, specific amounts of these ions play a role in the structure and function of the enzymes and bio-molecules inside or outside the cell. Iron ion increases the activity of some matrix metalloproteinase (MMPs). MMPs enzymes play a pivotal role in proliferation and remodeling processes of wound healing (166). Despite the reported negative effect of localized increase of Fe ion in chronic venous wounds (167, 168), the iron-chelating compounds accelerate the wound healing (169, 170). Lack of enough evidence for an appropriate range of these ions in the wound area requires more consideration in this field of research.

Many topical formulations recommended in TPM are a combination of several components. The composition and dose of ingredients have a great variety, preparing methods of traditional formulations do not match the current standards and criteria, and their preparation is time-consuming and difficult. Thus, the preparation method and dosing of formulations were not mentioned in the current study. However, new evidence for traditional combinations is increasingly available by different types of searches.

### Conclusion

Although some managements of burn wounds in TPM resources have been confirmed by new evidence, the efficacy of a plenty of materials and procedures needs to be methodically evaluated. Hence, it could introduce new clues for future studies.

### Conflict of interest

The authors declare that there are no conflicts of interest.

### References

1. World Health Organization. The global burden of disease. [cited 2020 Oct 25] Available from: [https://www.who.int/healthinfo/global\\_burden\\_disease/GBD\\_report\\_2004update\\_full.pdf](https://www.who.int/healthinfo/global_burden_disease/GBD_report_2004update_full.pdf).
2. Brusselsaers N, Monstrey S, Vogelaers D, Hoste E, Blot S. Severe burn injury in Europe: a systematic review of the incidence, etiology, morbidity, and mortality. *Crit Care* 2010; 14(5):R188. doi: 10.1186/cc9300.
3. Rowan MP, Cancio LC, Elster EA, Burmeister DM, Rose LF, Natesan S, et al. Burn wound healing and treatment: review and advancements. *Crit Care* 2015; 19:243. doi: 10.1186/s13054-015-0961-2.
4. Duke JM, Randall SM, Wood FM, Boyd JH, Fear MW. Burns and long-term infectious disease morbidity: A population-based study. *Burns* 2017; 43(2):273-81. doi: 10.1016/j.burns.2016.10.020.
5. Mehdizadeh A, Akbarian A, Samareh PP, Tavajjohi S, MacKay R, Alaghebandan R, et al. Epidemiology of burn injuries during pregnancy in Tehran, Iran. *Ann Burns Fire Disasters* 2002; 15(4):163-9.
6. Rae L, Fidler P, Gibran N. The physiologic basis of burn shock and the need for aggressive fluid resuscitation. *Crit Care Clin* 2016; 32(4):491-505. doi: 10.1016/j.ccc.2016.06.001.
7. Salas Campos L, Fernández Mansilla M, Martínez de la Chica AM. Topical chemotherapy for the treatment of burns. *Rev Enferm* 2005; 28(5):67-70. [In Spanish].
8. Hassanzadeh G, Hajmanouchehri F, Roi AB, Hassanzadeh N, Shafigh N, Barzroudpour M, et al. Comparing effects of silver sulfadiazine, sucralfate and brassica oleracea extract on burn wound healing. *Life Science Journal* 2013; 10(SUPPL):104-13.
9. Aziz Z, Abu SF, Chong NJ. A systematic review of silver-containing dressings and topical silver agents (used with dressings) for burn wounds. *Burns* 2012; 38(3):307-18. doi: 10.1016/j.burns.2011.09.020.
10. Oryan A, Alemzadeh E, Moshiri A. Burn wound healing: present concepts, treatment strategies and future directions. *J Wound Care* 2017; 26(1):5-19. doi: 10.12968/jowc.2017.26.1.5.
11. Pereira RF, Bartolo PJ. Traditional therapies for skin wound healing. *Adv Wound Care (New Rochelle)* 2016; 5(5):208-29. doi: 10.1089/wound.2013.0506.

12. Kopp J, Wang GY, Horch RE, Pallua N, Ge SD. Ancient traditional Chinese medicine in burn treatment: a historical review. *Burns* 2003; 29(5):473-8. doi: 10.1016/s0305-4179(03)00053-6.
13. Das U, Behera SS, Pramanik K. Ethno-Herbal-medico in wound repair: an incisive review. *Phytother Res* 2017; 31(4):579-90. doi: 10.1002/ptr.5786.
14. Bahramsoltani R, Farzaei MH, Abdolghaffari AH, Rahimi R, Samadi N, Heidari M, et al. Evaluation of phytochemicals, antioxidant and burn wound healing activities of *Cucurbita moschata* Duchesne fruit peel. *Iran J Basic Med Sci* 2017; 20(7):798-805. doi: 10.22038/IJBMS.2017.9015.
15. Mehrabani M, Seyyedkazemi SM, Nematollahi MH, Jafari E, Mehrabani M, Mehdipour M, et al. Accelerated burn wound closure in mice with a new formula based on traditional medicine. *Iran Red Crescent Med J* 2016; 18(11):e26613. doi: 10.5812/ircmj.26613.
16. Mehrbani M, Choopani R, Fekri A, Mehrabani M, Mosaddegh M, Mehrabani M. The efficacy of whey associated with dodder seed extract on moderate-to-severe atopic dermatitis in adults: A randomized, double-blind, placebo-controlled clinical trial. *J Ethnopharmacol* 2015; 172:325-32. doi: 10.1016/j.jep.2015.07.003.
17. Ebrahimipour N, Mehrabani M, Iranpour M, Kordestani Z, Mehrabani M, Nematollahi MH, et al. The efficacy of a traditional medicine preparation on second-degree burn wounds in rats. *J Ethnopharmacol* 2020; 252:112570. doi: 10.1016/j.jep.2020.112570.
18. Rameshk M, Sharififar F, Mehrabani M, Pardakhty A, Farsinejad A, Mehrabani M. Proliferation and in vitro wound healing effects of the microniosomes containing narcissus tazetta L. Bulb Extract on Primary Human Fibroblasts (HDFs). *Daru* 2018; 26(1):31-42. doi: 10.1007/s40199-018-0211-7.
19. Choopani R, Mehrbani M, Fekri A, Mehrabani M. Treatment of atopic dermatitis from the perspective of traditional persian medicine: presentation of a novel therapeutic approach. *J Evid Based Complementary Altern Med* 2017; 22(1):5-11. doi: 10.1177/2156587215598610.
20. Ibn Sina. *Canon of Medicine*. Beirut: Ehya al-Toras al-Arabi Press; 2010. [In Arabic].
21. Gorgani Z. *Zakhireh-e Kharazmshahi*. Qom: Rehabilitation Institutes Of Natural Medicine; 2012. [In Persian].
22. al-Razi MZ. *Al-Hawi*. Tehran: Academy of Medical Sciences; 2005. [In Arabic].
23. al-Razi MZ. *Al-Mansouri fi al-Tibb*. Tehran: Tehran Medical University; 1999. [In Arabic].
24. Aghili Khorasani MH. *Makhzan al-Advieh*. Tehran: Rah-e Kamal Publications; 2009. [In Persian].
25. Kermani M. *Daghayegh al-Alaj*. Kerman: Saadat Press; 1983. [In Persian].
26. al-Razi MZ. *Bar-us-sa'ah*. Tehran: Iran University of Medical Sciences; 2005. [In Arabic].
27. Chashty MA. *Romuz-e Azam*. Tehran: Iran University of Medical Sciences; 2004. [In Persian].
28. Kordafshari G, Ardakani MR, Keshavarz M, Esfahani MM, Nazem E, Moghimi M, et al. The role of phlebotomy (fasd) and wet cupping (hijamat) to manage dizziness and vertigo from the viewpoint of persian medicine. *J Evid Based Complementary Altern Med* 2017; 22(3):369-73. doi: 10.1177/2156587216672757.
29. Arzani MA. *Mofareh al-Gholub*. Lahoor: Salim Lahoor; 1915. [In Persian].
30. Avijgan M, Kamran A, Abedini A. Effectiveness of aloe vera gel in chronic ulcers in comparison with conventional treatments. *Iran J Med Sci* 2016; 41(3 Suppl):S30.
31. Goudarzi M, Fazeli M, Azad M, Seyedjavadi SS, Mousavi R. Aloe vera gel: effective therapeutic agent against multidrug-resistant *Pseudomonas aeruginosa* isolates recovered from burn wound infections. *Chemother Res Pract* 2015; 2015:639806. doi: 10.1155/2015/639806.
32. Akhoondinasab MR, Akhoondinasab M, Saberi M. Comparison of healing effect of aloe vera extract and silver sulfadiazine in burn injuries in experimental rat model. *World J Plast Surg* 2014; 3(1):29-34.
33. Liu H, Lin S, Xiao D, Zheng X, Gu Y, Guo S. Evaluation of the wound healing potential of resina draconis (*dracaena cochinchinensis*) in animal models. *Evid Based Complement Alternat Med* 2013; 2013:709865. doi: 10.1155/2013/709865.
34. Okuma CH, Andrade TA, Caetano GF, Finci LI, Maciel NR, Topan JF, et al. Development of lamellar gel phase emulsion containing marigold oil (*Calendula officinalis*) as a potential modern wound dressing. *Eur J Pharm Sci* 2015; 71:62-72. doi: 10.1016/j.ejps.2015.01.016.

35. Sengupta R. Combined wound healing activity of *Calendula officinalis* and Basil leaves. *J Pharmacog Phytochem* 2017; 6(1):173-76.
36. Parente LM, Lino Júnior Rde S, Tresvenzol LM, Vinaud MC, de Paula JR, Paulo NM, et al. Wound healing and anti-inflammatory effect in animal models of *Calendula officinalis* L. growing in Brazil. *Evid Based Complement Alternat Med* 2012; 2012:375671. doi: 10.1155/2012/375671.
37. Millán D, Jiménez RA, Nieto LE, Linero I, Laverde M, Fontanilla MR. Preclinical evaluation of collagen type I scaffolds, including gelatin-collagen microparticles and loaded with a hydroglycolic *Calendula officinalis* extract in a lagomorph model of full-thickness skin wound. *Drug Deliv Transl Res* 2016; 6(1):57-66. doi: 10.1007/s13346-015-0265-8.
38. Shafeie N, Tabatabai Naini A, Kargar Jahromi H. Comparison of different concentrations of *Calendula officinalis* gel on cutaneous wound healing. *Biomedical and Pharmacology Journal* 2015; 8(2):979-92. doi: 10.13005/bpj/850.
39. Buzzi M, de Freitas F, de Barros Winter M. Therapeutic effectiveness of a *Calendula officinalis* extract in venous leg ulcer healing. *J Wound Care* 2016; 25(12):732-9. doi: 10.12968/jowc.2016.25.12.732.
40. Parente LM, Andrade MA, Brito LA, Moura VM, Miguel MP, Lino-Junior Rde S, et al. Angiogenic activity of *Calendula officinalis* flowers L. in rats. *Acta Cir Bras* 2011; 26(1):19-24. doi: 10.1590/s0102-86502011000100005.
41. Sutar I, Kupeli Akkol E, Keles H, Yesilada E, Sarker SD, Baykal T. Comparative evaluation of traditional prescriptions from *Cichorium intybus* L. for wound healing: stepwise isolation of an active component by in vivo bioassay and its mode of activity. *J Ethnopharmacol* 2012; 143(1):299-309. doi: 10.1016/j.jep.2012.06.036.
42. Heidari M, Bahramsoltani R, Abdolghaffari AH, Rahimi R, Esfandyari M, Baeri M, et al. Efficacy of topical application of standardized extract of *Tragopogon graminifolius* in the healing process of experimental burn wounds. *J Tradit Complement Med* 2019; 9(1):54-9. doi: 10.1016/j.jtcme.2018.02.002.
43. Ashkani-Esfahani S, Imanieh MH, Khoshneviszadeh M, Meshksar A, Noorafshan A, Geramizadeh B, et al. The healing effect of *Arnebia euchroma* in second degree burn wounds in rat as an animal model. *Iran Red Crescent Med J* 2012; 14(2):70-4.
44. Nasiri E, Hosseinimehr SJ, Azadbakht M, Akbari J, Enayati-Fard R, Azizi S, et al. The healing effect of *Arnebia euchroma* ointment versus silver sulfadiazine on burn wounds in rat. *World J Plast Surg* 2015; 4(2):134-44.
45. Mohsenikia M, Khakpour S, Azizian Z, Ashkani-Esfahani S, Razavipour ST, Toghiani P. Wound healing effect of *Arnebia euchroma* gel on excisional wounds in rats. *Adv Biomed Res* 2017; 6:2. doi: 10.4103/2277-9175.199260.
46. Aliasl J, Barikbin B, Khoshzaban F, Naseri M, Sedaghat R, Kamalinejad M, et al. Effect of *Arnebia euchroma* ointment on post-laser wound healing in rats. *J Cosmet Laser Ther* 2015; 17(1):41-5. doi: 10.3109/14764172.2014.968583.
47. Ghasemi Pirbalouti A, Yousefi M, Nazari H, Karimi I, Koohpayeh A. Evaluation of burn healing properties of *Arnebia euchroma* and *Malva sylvestris*. *Electronic Journal of Biology* 2009; 5(3):62-6.
48. Sarandy MM, Novaes RD, da Matta SL, Mezcencio JM, da Silva MB, Zanuncio JC, et al. Ointment of *Brassica oleracea* var. capitata matures the extracellular matrix in skin wounds of wistar rats. *Evid Based Complement Alternat Med* 2015; 2015:919342. doi: 10.1155/2015/919342.
49. Rebolla A, Arisawa EA, Barja PR, Posso MB, Carvalho Cda S. Effect of *Brassica oleracea* in rats skin wound healing. *Acta Cir Bras* 2013; 28(9):664-9. doi: 10.1590/s0102-86502013000900007.
50. Goncalves RV, Sarandy MM, da Matta SL, Novaes RD, Pinto MV. Comparative study of the effects of laser photobiomodulation and extract of *Brassica oleracea* on skin wounds in wistar rats: a histomorphometric study. *Pathology - Research and Practice* 2013; 209(10):648-53. doi: 10.1016/j.prp.2013.07.006.
51. Hou Q, He WJ, Chen L, Hao HJ, Liu JJ, Dong L, et al. Effects of the four-herb compound anbp on wound healing promotion in diabetic mice. *Int J Low Extrem Wounds* 2015; 14(4):335-42. doi: 10.1177/1534734615575244.
52. Davis SC, Mertz PM. Determining the effect of an oak bark formulation on methicillin-resistant *Staphylococcus aureus* and wound healing in porcine wound models. *Ostomy Wound Manage* 2008; 54(10):16-8, 20, 22-5.
53. Ali M, Abdel Motaal A, Ahmed MA, Alsayari A, El-Gazayerly ON. An in vivo study of *Hypericum perforatum* in a niosomal topical drug delivery system. *Drug Deliv*. 2018;

- 25(1):417-25. doi: 10.1080/10717544.2018.1431977.
54. Farsak M, Özdağlı G, Özmüş D, Çömelekoğlu Ü, Yalın S, Bozdoğan Arpacı R, et al. Effects of *Hypericum perforatum* on an experimentally induced diabetic wound in a rat model. *Wounds* 2017; 29(2):E10-7.
  55. Yadollah-Damavandi S, Chavoshi-Nejad M, Jangholi E, Nekouyian N, Hosseini S, Seifae A, et al. Topical *hypericum perforatum* improves tissue regeneration in full-thickness excisional wounds in diabetic rat model. *Evidence-Based Complementary and Alternative Medicine* 2015; 2015: 245328. doi: 10.1155/2015/245328.
  56. Kíyan S, Uyanıkgil Y, Altuncı YA, Cavusoglu T, Cetin Uyanıkgil EO, Karabey F. Investigation of acute effects of *Hypericum perforatum* (St. John's Wort-Kantaron) treatment in experimental thermal burns and comparison with silver sulfadiazine treatment. *Ulus Travma Acil Cerrahi Derg* 2015; 21(5):323-36. doi: 10.5505/tjtes.2015.
  57. Prisacaru AI, Andritoiu CV, Andriescu C, Havarneanu EC, Popa M, Motoc AG, et al. Evaluation of the wound-healing effect of a novel *Hypericum perforatum* ointment in skin injury. *Rom J Morphol Embryol* 2013; 54(4):1053-9.
  58. Mukherjee PK, Verpoorte R, Suresh B. Evaluation of in-vivo wound healing activity of *Hypericum patulum* (Family: hypericaceae) leaf extract on different wound model in rats. *J Ethnopharmacol* 2000; 70(3):315-21. doi: 10.1016/s0378-8741(99)00172-5.
  59. Khorasani G, Hosseinimehr SJ, Zamani P, Ghasemi M, Ahmadi A. The effect of saffron (*Crocus sativus*) extract for healing of second-degree burn wounds in rats. *Keio J Med* 2008; 57(4):190-5. doi: 10.2302/kjm.57.190.
  60. Solanki R, Mathur V, Purohit SK, Mathur M. Evaluation of wound healing activity of ethanolic extract of *Ocimum basilicum* and *Aegle marmelos* leaves in male albino rats. *International Journal of Drug Research and Technology* 2017; 2(4).
  61. Shetty S, Udupa S, Udupa L. Evaluation of antioxidant and wound healing effects of alcoholic and aqueous extract of *ocimum sanctum* linn in rats. *Evid Based Complement Alternat Med* 2008; 5(1):95-101. doi: 10.1093/ecam/nem004.
  62. Osuagwu FC, Oladejo OW, Imosemi IO, Adewoyin BA, Aiku A, Ekpo OE, et al. Wound healing activities of methanolic extracts of *Ocimum gratissimum* leaf in Wistar rats - a preliminary study. *Afr J Med Med Sci* 2004; 33(1):23-6.
  63. Orafidiya LO, Agbani EO, Abereoje OA, Awe T, Abudu A, Fakoya FA. An investigation into the wound-healing properties of essential oil of *Ocimum gratissimum* linn. *J Wound Care* 2003; 12(9):331-4. doi: 10.12968/jowc.2003.12.9.26537.
  64. Vaghardoost R, Mousavi Majd SG, Tebyanian H, Babavalian H, Malaei L, Niazi M, et al. The healing effect of sesame oil, camphor and honey on second degree burn wounds in rat. *World J Plast Surg* 2018; 7(1):67-71.
  65. Singh S, Sharma N. Evaluation of wound healing activity of *Acacia auriculiformis* A. Cunn. Stem bark. *Asain J Pharm Clin Res* 2014; 7(2):204-7.
  66. Suriyamoorthy S, Subramaniam K, Durai SJ, Wahaab F, Chitraselvi RP. Evaluation of wound healing activity of *Acacia caesia* in rats. *Wound Medicine* 2014; 7:1-7. doi: 10.1016/j.wndm.2015.03.001.
  67. Castro B, Palomares T, Azcoitia I, Bastida F, del Olmo M, Soldevilla JJ, et al. Development and preclinical evaluation of a new galactomannan-based dressing with antioxidant properties for wound healing. *Histol Histopathol* 2015; 30(12):1499-512. doi: 10.14670/HH-11-646.
  68. Sumitra M, Manikandan P, Suguna L, Cehittar G. Study of dermal wound healing activity of *Trigonella foenum graecum* seeds in rats. *J Clin Biochem Nutr* 2000; 28(2):59-67.
  69. Ktari N, Trabelsi I, Bardaa S, Triki M, Bkhairia I, Salem RBS-B, et al. Antioxidant and hemolytic activities, and effects in rat cutaneous wound healing of a novel polysaccharide from fenugreek (*Trigonella foenum-graecum*) seeds. *Int J Biol Macromol* 2017; 95:625-34. doi: 10.1016/j.ijbiomac.2016.11.091.
  70. Mustafa MR, Mahmood AA, Salmah I. Effects of *Trigonella foenum-graecum* seed extract in combination with honey on experimental wound healing in rats. *International Journal of Molecular Medicine and Advance Science* 2005; 1(1):29-33.
  71. Farahpour MR, Aghaei M. Assessment of the effect of co-administration of aloe vera gel and fenugreek seed hydroethanolic extract on the improvement of full-thickness excisional skin wound healing in diabetic mice. *Veterinary Clinical Pathology (Veterinary Journal Tabriz)* 2016; 9(36):285-95. [In Persian].

72. Moalla Rekik D, Ben Khedir S, Ksouda Moalla K, Kammoun NG, Rebai T, Sahnoun Z. Evaluation of wound healing properties of grape seed, sesame, and fenugreek oils. *Evidence-Based Complementary and Alternative Medicine* 2016; 2016: 7965689. doi: 10.1155/2016/7965689.
73. Farahpour MR, Aghaei M. Effect fenugreek seeds hydroethanolic extract on a full-thickness wound healing in streptozotocin-induced diabetic mice. *Journal of Comparative Pathobiology Iran* 2016; 13(54):1979-86.
74. Muhammed DO, Salih NA. Effect of application of Fenugreek (*Trigonella foenum-graecum*) on skin wound healing in rabbits. *AL-Qadisiyah Journal of Veterinary Medicine Sciences* 2012; 11(2):86-93.
75. Sudheendra AT, Shenoy R, Taranalli AD, MCOPS M. Evaluation of flavonoid fraction of *Trigonella foenum-graecum* for its wound healing potential. *Pharmacologyonline* 2009; 1:1154-62.
76. Selvaraj S, Fathima NN. Fenugreek incorporated silk fibroin nanofibers-a potential antioxidant scaffold for enhanced wound healing. *ACS Appl Mater Interfaces* 2017; 9(7):5916-26. doi: 10.1021/acsami.6b16306.
77. Farahpour MR, Taghikhani H, Habibi M, Zandieh MA. Wound healing activity of flaxseed *Linum usitatissimum* L. in rats. *Afr J Pharm Pharmacol* 2011; 5(21):2386-9.
78. de Souza Franco E, de Aquino CM, de Medeiros PL, Evêncio LB, da Silva Góes AJ, de Souza Maia MB. Effect of a semisolid formulation of *Linum usitatissimum* L.(Linseed) oil on the repair of skin wounds. *Evid Based Complement Alternat Med* 2012; 2012:270752. doi: 10.1155/2012/270752.
79. Datta HS, Mitra SK, Patwardhan B. Wound healing activity of topical application forms based on ayurveda. *Evid Based Complement Alternat Med* 2011; 2011:134378. doi: 10.1093/ecam/nep015.
80. Jridi M, Sellimi S, Lassoued KB, Beltaief S, Souissi N, Mora L, et al. Wound healing activity of cuttlefish gelatin gels and films enriched by henna (*Lawsonia inermis*) extract. *Colloids Surf A Physicochem Eng Asp.* 2017; 512:71-9.
81. Shiravi AH, Alebooyeh M, Hojati V, Akbari H. The effect of extract of henna leaves (*Lawsonia inermis*) on skin wound healing in wistar rats. *Journal of Animal Biology* 2011; 3(4):45-51. [In Persian].
82. Nayak BS, Isitor G, Davis EM, Pillai GK. The evidence based wound healing activity of *Lawsonia inermis* Linn. *Phytother Res* 2007; 21(9):827-31. doi: 10.1002/ptr.2181.
83. Afshar M, Ravarian B, Zardast M, Moallem SA, Fard MH, Valavi M. Evaluation of cutaneous wound healing activity of *Malva sylvestris* aqueous extract in BALB/c mice. *Iran J Basic Med Sci* 2015; 18(6):616-22.
84. Pirbalouti AG, Azizi S, Koochpayeh A, Hamed B. Wound healing activity of *Malva sylvestris* and *Punica granatum* in alloxan-induced diabetic rats. *Acta Pol Pharm* 2010; 67(5):511-6.
85. Fahimi S, Abdollahi M, Mortazavi SA, Hajimehdipoor H, Abdolghaffari AH, Rezvanfar MA. Wound healing activity of a traditionally used poly herbal product in a burn wound model in rats. *Iran Red Crescent Med J* 2015; 17(9):e19960. doi: 10.5812/ircmj.19960.
86. Barua C, Talukdar A, Barua AG, Chakraborty A, Sarma RK, Bora RS. Evaluation of the wound healing activity of methanolic extract of *Azadirachta Indica* (Neem) and *Tinospora cordifolia* (Guduchi) in rats. *Pharmacologyonline* 2010; 1:70-7.
87. Singh A, Singh AK, Narayan G, Singh TB, Shukla VK. Effect of Neem oil and Haridra on non-healing wounds. *Ayu* 2014; 35(4):398-403. doi: 10.4103/0974-8520.158998.
88. Babu KS, Naik VK, Latha J, Prabhakar V. Wound healing activity of ethanolic extract of natural products (*Azadirachta indica* bark) in albino wister rats. *World J Pharm Pharm Sci* 2016; 5(6):1624-32.
89. Shukla R, Kashaw SK, Jain AP, Lodhi S. Fabrication of Apigenin loaded gellan gum-chitosan hydrogels (GGCH-HGs) for effective diabetic wound healing. *Int J Biol Macromol* 2016; 91:1110-9. doi: 10.1016/j.ijbiomac.2016.06.075.
90. Amutha K, Selvakumari U. Wound healing activity of methanolic stem extract of *Musa paradisiaca* Linn. (Banana) in Wistar albino rats. *Int Wound J* 2016; 13(5):763-7. doi: 10.1111/iwj.12371.
91. Ozcan O, Ipekci H, Alev B, Ustundag UV, Ak E, Sen A, et al. Protective effect of Myrtle (*Myrtus communis*) on burn induced skin injury. *Burns* 2019; 45(8):1856-63. doi: 10.1016/j.burns.2019.07.015.
92. Raeiszadeh M, Esmaeili-Tarzi M, Bahrapour-Juybari K, Nematollahi-Mahani SN, Pardakhty A, Nematollahi MH, et al. Evaluation the effect of *Myrtus communis* L. extract on several

- underlying mechanisms involved in wound healing: An in vitro study. *S Afr J Bot* 2018; 118:144-50. doi: 10.1016/j.sajb.2018.07.006.
93. Hashemipour MA, Lotfi S, Torabi M, Sharifi F, Ansari M, Ghassemi A, et al. evaluation of the effects of three plant species (*myrtus communis* L., *camellia sinensis* L., *zataria multiflora* Boiss.) on the healing process of intraoral ulcers in rats. *J Dent (Shiraz)* 2017; 18(2):127-35.
94. Zahmatkesh M, Manesh MJ, Babashahabi R. Effect of Olea ointment and Acetate Mafenide on burn wounds - A randomized clinical trial. *Iran J Nurs Midwifery Res* 2015; 20(5):599-603. doi: 10.4103/1735-9066.164507.
95. Koca U, Süntar I, Akkol EK, Yılmaz D, Alper M. Wound repair potential of *Olea europaea* L. leaf extracts revealed by in vivo experimental models and comparative evaluation of the extracts' antioxidant activity. *J Med Food* 2011; 14(1-2):140-6. doi: 10.1089/jmf.2010.0039.
96. Kazemi-Darabadi S, Akbari G, Jarolmasjed SH, Shahbazfar AA. A histopathologic study of effects of olive oil plus lime water on third-degree burn in mouse model. *Iranian Journal of Veterinary Surgery* 2017; 12(1):55-63. doi: 10.22034/ivsa.2017.50916.
97. Akbari G, Shahbazfar A, Kianifard D, Rezaei H, Shokrollahi S, Mohebi D. Microscopic study of the healing effects of the mixture of olive oil and lime water on the second degree burning in rats. *Journal of Ilam University of Medical Sciences* 2017; 24(6):169-77. doi: 10.18869/acadpub.sjimu.24.6.169.
98. Tumen I, Süntar I, Keleş H, Küpeli Akkol E. A therapeutic approach for wound healing by using essential oils of *Cupressus* and *Juniperus* species growing in Turkey. *Evid Based Complement Alternat Med* 2012; 2012:728281. doi: 10.1155/2012/728281.
99. Shenoy RR, Sudheendra AT, Nayak PG, Paul P, Kutty NG, Rao CM. Normal and delayed wound healing is improved by sesamol, an active constituent of *Sesamum indicum* (L.) in albino rats. *J Ethnopharmacol* 2011; 133(2):608-12. doi: 10.1016/j.jep.2010.10.045.
100. Kiran K, Asad M. Wound healing activity of *Sesamum indicum* L seed and oil in rats. *Indian J Exp Biol* 2008; 46(11):777-82.
101. Sharif M, Alizargar J, Sharif A. Evaluation of the wound healing activity of sesame oil extract in rats. *World Journal of Medical Sciences* 2013; 9(2):74-8. doi: 10.5829/idosi.wjms.2013.9.2.75195.
102. Tehrani S, Lotfi P, Tehrani S, Jangholi E, Aryan H, Aidun A. Healing effect of sesame ointment on second-degree burn wound in rats. *Galen Medical Journal* 2016; 5(2):56-62.
103. Bazzi P, Akbarzadeh S, Obeidi N, Noohpisheh MK, Daneshi A, Bargahi A. Using toasted barley in sesame oil mixture for non-surgical necrosis debridement of experimental burns in rat. *J Med Plant Res* 2014; 8(1):81-7. doi: 10.5897/JMPR2013.5183.
104. Amini M, Kherad M, Mehrabani D, Azarpira N, Panjehshahin MR, Tanideh N. Effect of plantago major on burn wound healing in rat. *J Appl Anim Res* 2010; 37(1):53-6. doi: 10.1080/09712119.2010.9707093.
105. Thomé RG, dos Santos HB, dos Santos FV, da Silva Oliveira RJ, de Camargos LF, Pereira MN, et al. Evaluation of healing wound and genotoxicity potentials from extracts hydroalcoholic of *Plantago major* and *Siparuna guianensis*. *Exp Biol Med (Maywood)* 2012; 237(12):1379-86. doi: 10.1258/ebm.2012.012139.
106. Patil BS, Mastiholimath VS, Kulkarni AR. Development and evaluation of psyllium seed husk polysaccharide based wound dressing films. *Orient Pharm Exp Med* 2011; 11:123. doi: 10.1007/s13596-011-0012-8.
107. Hemmati AA, Kalantari H, Jalali A, Rezai S, Zadeh HH. Healing effect of quince seed mucilage on T-2 toxin-induced dermal toxicity in rabbit. *Exp Toxicol Pathol* 2012; 64(3):181-6. doi: 10.1016/j.etp.2010.08.004.
108. Omidian M, Hemmati AA, Farajzade H, Houshmand G, Sattari A, Maryam K. Priority of 5% quince seed cream versus 1% phenytoin cream in the healing of skin ulcers: a randomized controlled trial. *Jundishapur J Nat Pharm Prod* 2015; 10(2): e24590. doi: 10.17795/jjnpp-24590.
109. Mansouri E, Hardani A, Afzalzadeh MR, Amir Zargar A, Meamar Z. Combined effects of retinoic acid and hydro-alcoholic extract of *rosa damascena* mill on wound in diabetic rats. *Iran J Pharm Res* 2016; 15(2):583-9.
110. d'Alessio PA, Mirshahi M, Bisson JF, Bene MC. Skin repair properties of d-Limonene and perillyl alcohol in murine models. *Antiinflamm Antiallergy Agents Med Chem* 2014; 13(1):29-35. doi: 10.2174/18715230113126660021.
111. Yusufoglu HS, Alqasoumi SI. Anti-inflammatory and wound healing activities of herbal gel containing an antioxidant *Tamarix aphylla* leaf extract. *International Journal of Pharmacology* 2011; 7(8):829-35. doi: 10.3923/ijp.2011.829.835.



112. Demling RH, Leslie DeSanti MD. The rate of re-epithelialization across meshed skin grafts is increased with exposure to silver. *Burns* 2002; 28(3):264-6. doi: 10.1016/s0305-4179(01)00119-x.
113. Kucan JO, Robson MC, Hegggers JP, Ko F. Comparison of silver sulfadiazine, povidone-iodine and physiologic saline in the treatment of chronic pressure ulcers. *J Am Geriatr Soc* 1981; 29(5):232-5. doi: 10.1111/j.1532-5415.1981.tb01773.x.
114. Luo GG, Huang RB, Chen JH, Lin LR, Zheng LS. Diversity of coordination architecture of silver(I) complexes with different 2-aminopyrimidyl derivatives: Effect of counter anions and ligands. *Polyhedron* 2008; 27(13):2791-8. doi: 10.1016/j.poly.2008.06.003.
115. Chen J, Han CM, Lin XW, Tang ZJ, Su SJ. Effect of silver nanoparticle dressing on second degree burn wound. *Zhonghua Wai Ke Za Zhi* 2006; 44(1):50-2. [In Chinese].
116. Lohsirawat V, Chuangsuwanich A. Comparison of the ionic silver-containing hydrofiber and paraffin gauze dressing on split-thickness skin graft donor sites. *Ann Plast Surg* 2009; 62(4):421-2. doi: 10.1097/SAP.0b013e31818a65e9.
117. Otsuka M, Hatakeyama H, Shikamura M, Otsuka K, Ito A. Therapeutic effects of transdermal systems containing zinc-related materials on thermal burn rats. *Biomed Mater Eng* 2015; 25(2):143-56. doi: 10.3233/BME-151265.
118. Kaufman KL, Mann FA, Kim DY, Lee S, Yoon HY. Evaluation of the effects of topical zinc gluconate in wound healing. *Vet Surg* 2014; 43(8):972-82. doi: 10.1111/j.1532-950X.2014.12243.x.
119. Arslan K, Karahan O, Okuş A, Unlü Y, Eryılmaz MA, Ay S, et al. Comparison of topical zinc oxide and silver sulfadiazine in burn wounds: an experimental study. *Ulus Travma Acil Cerrahi Derg* 2012; 18(5):376-83. doi: 10.5505/tjtes.2012.45381.
120. Thompson CB, Wiemken TL, Brown TS. Effect of postoperative dressing on excisions performed on the leg: a comparison between zinc oxide compression dressings versus standard wound care. *Dermatol Surg* 2017; 43(11):1379-84. doi: 10.1097/DSS.0000000000001209.
121. Mulder GD, Patt LM, Sanders L, Rosenstock J, Altman MI, Hanley ME, et al. Enhanced healing of ulcers in patients with diabetes by topical treatment with glycyl-l-histidyl-l-lysine copper. *Wound Repair Regen* 1994; 2(4):259-69. doi: 10.1046/j.1524-475X.1994.20406.x.
122. Borkow G, Gabbay J, Dardik R, Eidelman AI, Lavie Y, Grunfeld Y, et al. Molecular mechanisms of enhanced wound healing by copper oxide-impregnated dressings. *Wound Repair Regen* 2010; 18(2):266-75. doi: 10.1111/j.1524-475X.2010.00573.x.
123. Malik KI, Malik MA, Aslam A. Honey compared with silver sulphadiazine in the treatment of superficial partial-thickness burns. *Int Wound J* 2010; 7(5):413-7. doi: 10.1111/j.1742-481X.2010.00717.x.
124. Lavaf M, Simbar M, Mojab F, Alavi Majd H, Samimi M. Comparison of honey and phenytoin (PHT) cream effects on intensity of pain and episiotomy wound healing in nulliparous women. *J Complement Integr Med* 2017; 15(1). doi: 10.1515/jcim-2016-0139.
125. Goharshenasan P, Amini S, Atria A, Abtahi H, Khorasani G. Topical application of honey on surgical wounds: a randomized clinical trial. *Forsch Komplementmed* 2016; 23(1):12-5. doi: 10.1159/000441994.
126. Takzaree N, Hassanzadeh G, Rouini MR, Manayi A, Hadjiakhondi A, Majidi Zolbin M. Evaluation of the effects of local application of thyme honey in open cutaneous wound healing. *Iran J Public Health* 2017; 46(4):545-51.
127. Mphande AN, Killowe C, Phalira S, Jones HW, Harrison WJ. Effects of honey and sugar dressings on wound healing. *J Wound Care*. 2007; 16(7):317-9. doi: 10.12968/jowc.2007.16.7.27053.
128. Paydar S, Akrami M, Dehghanian A, Alavi Moghadam R, Heidarpour M, Bahari Khoob A, et al. A comparison of the effects of alpha and medical-grade honey ointments on cutaneous wound healing in rats. *J Pharm (Cairo)* 2016; 2016:9613908. doi: 10.1155/2016/9613908.
129. Alizadeh AM, Sohanaki H, Khaniki M, Mohaghheghi MA, Ghmami G, Mosavi M. The effect of teucrium polium honey on the wound healing and tensile strength in rat. *Iran J Basic Med Sci* 2011; 14(6):499-505.
130. Chamani G, Zarei MR, Mehrabani M, Mehdavinezhad A, Vahabian M, Ahmadi-Motamayel F. Evaluation of Honey as a Topical Therapy for Intraoral Wound Healing in Rats. *Wounds* 2017; 29(3):80-6.
131. Ghaderi R, Afshar M. Topical application of honey for treatment of skin wound in mice. *Iran J Med Sci* 2004; 29(4):185-8.
132. Iftikhar F, Arshad M, Rasheed F, Amraiz D, Anwar P, Gulfraz M. Effects of acacia honey on

- wound healing in various rat models. *Phytother Res* 2010; 24(4):583-6. doi: 10.1002/ptr.2990.
133. Baghel PS, Shukla S, Mathur RK, Randa R. A comparative study to evaluate the effect of honey dressing and silver sulfadiazene dressing on wound healing in burn patients. *Indian J Plast Surg* 2009; 42(2):176-81. doi: 10.4103/0970-0358.59276.
134. Sirousazar M, Jahani-Javanmardi A, Kheiri F, Hassan ZM. In vitro and in vivo assays on egg white/polyvinyl alcohol/clay nanocomposite hydrogel wound dressings. *J Biomater Sci Polym Ed* 2016; 27(16):1569-83. doi: 10.1080/09205063.2016.1218210.
135. Fu H, Wang XJ, Lu HF, Xu QW. Experimental study on beeswax ointment in diabetic rabbit's model of wound healing. *Journal of Dalian Medical University* 2007; 4.
136. Kumari P, Chahar MK, Veerapur VP, Spandana G, Thippeswamy BS, Badami S. Spider web ointment: A traditional based approach in Cutaneous wound healing. *Indian Journal of Traditional Knowledge* 2013; 12(4):657-63.
137. Bhat ZA. Traditional medicines in drug discovery. *J Pharm Res* 2012; 5(5):2457-9.
138. Campelo AP, Campelo MW, Britto GA, Ayala AP, Guimarães SB, Vasconcelos PR. An optimized animal model for partial and total skin thickness burns studies. *Acta Cir Bras* 2011; 26(Suppl 1):38-42. doi: 10.1590/s0102-86502011000700008.
139. Carvalho AR Jr, Diniz RM, Suarez MA, Figueiredo CS, Zigmignan A, Grisotto MA, et al. Use of some asteraceae plants for the treatment of wounds: from ethnopharmacological studies to scientific evidences. *Front Pharmacol* 2018; 9:784. doi: 10.3389/fphar.2018.00784.
140. Denisow-Pietrzyk M, Pietrzyk Ł, Denisow B. Asteraceae species as potential environmental factors of allergy. *Environ Sci Pollut Res Int* 2019; 26(7):6290-300. doi: 10.1007/s11356-019-04146-w.
141. Salekzamani S, Ebrahimi-Mameghani M, Rezazadeh K. The antioxidant activity of artichoke (*Cynara scolymus*): A systematic review and meta-analysis of animal studies. *Phytother Res* 2019; 33(1):55-71. doi: 10.1002/ptr.6213.
142. Ben Salem M, Affes H, Dhoubi R, Charfi S, Turki M, Hammami S, et al. Effect of Artichoke (*cynara scolymus*) on cardiac markers, lipid profile and antioxidants levels in tissue of HFD-induced obesity. *Arch Physiol Biochem* 2019; 1-11. doi: 10.1080/13813455.2019.1670213.
143. Biel W, Witkiewicz R, Piątkowska E, Podsiadło C. Proximate composition, minerals and antioxidant activity of artichoke leaf extracts. *Biol Trace Elem Res* 2020; 194(2):589-95. doi: 10.1007/s12011-019-01806-3.
144. Pareek A, Suthar M, Rathore GS, Bansal V. Feverfew (*Tanacetum parthenium* L.): A systematic review. *Pharmacogn Rev* 2011; 5(9):103-10. doi: 10.4103/0973-7847.79105.
145. Bouziri A, Hamdi A, Menif K, Ben Jaballah N. Hepatorenal injury induced by cutaneous application of *Atractylis gummifera* L. *Clin Toxicol (Phila)* 2010; 48(7):752-4. doi: 10.3109/15563650.2010.498379.
146. Doulah A, Neisi N, Zekavati R, Farjam M. Antibacterial, antifungal and antioxidant activity of four species from *Arnebia* genus growing wild in Iran. *Iranian Journal of Science & Technology* 2014; 38(2):159-64.
147. Tasset-Cuevas I, Fernández-Bedmar Z, Lozano-Baena MD, Campos-Sánchez J, de Haro-Bailón A, Muñoz-Serrano A, et al. Protective effect of borage seed oil and gamma linolenic acid on DNA: in vivo and in vitro studies. *PLoS One* 2013; 8(2):e56986. doi: 10.1371/journal.pone.0056986.
148. Silva JR, Burger B, Kuhl CM, Candreva T, Dos Anjos MB, Rodrigues HG. Wound healing and omega-6 fatty acids: from inflammation to repair. *Mediators Inflamm* 2018; 2018:2503950. doi: 10.1155/2018/2503950.
149. Nogueira AO, Oliveira YI, Adjafre BL, de Moraes ME, Aragao GF. Pharmacological effects of the isomeric mixture of alpha and beta amyryn from *Protium heptaphyllum*: a literature review. *Fundam Clin Pharmacol* 2019; 33(1):4-12. doi: 10.1111/fcp.12402.
150. Zarshenas MM, Arabzadeh A, Ajdari Tafti M, Kordafshari G, Zargarani A, Mohagheghzadeh A. Application of herbal exudates in traditional Persian medicine. *Galen Medical Journal* 2012; 1(2):78-83.
151. Amen YM, Marzouk AM, Zaghoul MG, Afifi MS. The genus *Machaerium* (Fabaceae): taxonomy, phytochemistry, traditional uses and biological activities. *Nat Prod Res* 2015; 29(15):1388-405. doi: 10.1080/14786419.2014.1003062.
152. Wong VW, Martindale RG, Longaker MT, Gurtner GC. From germ theory to germ therapy: skin microbiota, chronic wounds, and probiotics. *Plast Reconstr Surg* 2013; 132(5):854e-61e. doi: 10.1097/PRS.0b013e3182a3c11e.

153. Baquerizo Nole KL, Yim E, Keri JE. Probiotics and prebiotics in dermatology. *J Am Acad Dermatol* 2014; 71(4):814-21. doi: 10.1016/j.jaad.2014.04.050.
154. Peral MC, Huaman Martinez MA, Valdez JC. Bacteriotherapy with *Lactobacillus plantarum* in burns. *Int Wound J* 2009; 6(1):73-81. doi: 10.1111/j.1742-481X.2008.00577.x.
155. Valdez JC, Peral MC, Rachid M, Santana M, Perdigon G. Interference of *Lactobacillus plantarum* with *Pseudomonas aeruginosa* in vitro and in infected burns: the potential use of probiotics in wound treatment. *Clin Microbiol Infect* 2005; 11(6):472-9. doi: 10.1111/j.1469-0691.2005.01142.x.
156. Jebur MS. Therapeutic efficacy of *Lactobacillus acidophilus* against bacterial isolates from burn wounds. *N Am J Med Sci* 2010; 2(12):586-91. doi: 10.4297/najms.2010.2586.
157. Chatterjee D. A controlled comparative study of the use of porcine xenograft in the treatment of partial thickness skin loss in an occupational health centre. *Curr Med Res Opin* 1978; 5(9):726-33. doi: 10.1185/03007997809110213.
158. Voineskos SH, Ayeni OA, McKnight L, Thoma A. Systematic review of skin graft donor-site dressings. *Plast Reconstr Surg* 2009; 124(1):298-306. doi: 10.1097/PRS.0b013e3181a8072f.
159. Ruszczak Z. Effect of collagen matrices on dermal wound healing. *Adv Drug Deliv Rev* 2003; 55(12):1595-611. doi: 10.1016/j.addr.2003.08.003.
160. Shevchenko RV, Sibbons PD, Sharpe JR, James SE. Use of a novel porcine collagen paste as a dermal substitute in full-thickness wounds. *Wound Repair Regen* 2008; 16(2):198-207. doi: 10.1111/j.1524-475X.2008.00360.x.
161. Chashty MA. *Great Elixir*. Tehran: Iran University of Medical Sciences (Republished by Institute of Medical History, Islamic and Complementary Medicine); 2005. [in Persian].
162. Heyneman A, Hoeksema H, Vandekerckhove D, Pirayesh A, Monstrey S. The role of silver sulphadiazine in the conservative treatment of partial thickness burn wounds: A systematic review. *Burns* 2016; 42(7):1377-86. doi: 10.1016/j.burns.2016.03.029.
163. You C, Li Q, Wang X, Wu P, Ho JK, Jin R, et al. Silver nanoparticle loaded collagen/chitosan scaffolds promote wound healing via regulating fibroblast migration and macrophage activation. *Sci Rep* 2017; 7(1):10489. doi: 10.1038/s41598-017-10481-0.
164. Pati R, Mehta RK, Mohanty S, Padhi A, Sengupta M, Vaseeharan B, et al. Topical application of zinc oxide nanoparticles reduces bacterial skin infection in mice and exhibits antibacterial activity by inducing oxidative stress response and cell membrane disintegration in macrophages. *Nanomedicine* 2014; 10(6):1195-208. doi: 10.1016/j.nano.2014.02.012.
165. Lansdown AB, Mirastschijski U, Stubbs N, Scanlon E, Ågren MS. Zinc in wound healing: Theoretical, experimental, and clinical aspects. *Wound Repair Regen* 2007; 15(1):2-16. doi: 10.1111/j.1524-475X.2006.00179.x.
166. Zamboni P, Scapoli G, Lanzara V, Izzo M, Fortini P, Legnaro R, et al. Serum iron and matrix metalloproteinase-9 variations in limbs affected by chronic venous disease and venous leg ulcers. *Dermatol Surg* 2005; 31(6):644-9. doi: 10.1111/j.1524-4725.2005.31611.
167. Zamboni P. The big idea: iron-dependent inflammation in venous disease and proposed parallels in multiple sclerosis. *J R Soc Med* 2006; 99(11):589-93. doi: 10.1258/jrsm.99.11.589.
168. Koh TJ, DiPietro LA. Inflammation and wound healing: the role of the macrophage. *Expert Rev Mol Med* 2011; 13:e23. doi: 10.1017/S1462399411001943.
169. Mohammadpour M, Behjati M, Sadeghi A, Fassihi A. Wound healing by topical application of antioxidant iron chelators: kojic acid and deferiprone. *Int Wound J* 2013; 10(3):260-4. doi: 10.1111/j.1742-481X.2012.00971.x.
170. Mehrabani M, Najafi M, Kamarul T, Mansouri K, Iranpour M, Nematollahi MH, et al. Deferoxamine preconditioning to restore impaired HIF-1 $\alpha$ -mediated angiogenic mechanisms in adipose-derived stem cells from STZ-induced type 1 diabetic rats. *Cell Prolif* 2015; 48(5):532-49. doi: 10.1111/cpr.12209.