



Impact of Medicine Ball Training on Amplitude of Electromyography Activity in Back Pain Patients During Gait

Ehsan Fakhri Mirzang¹ , AmirAli Jafarnezhadgero^{2*} , Lotfali Bolboli³

¹Department of Physical Education and Sport Sciences, Faculty of Humanities and Social Sciences, University of Kurdistan, Sanandaj, Iran

²Department of Sport Biomechanics, Faculty of Educational Science and Psychology, University of Mohaghegh Ardabili, Ardabil, Iran

³Department of Sport Physiology, Faculty of Educational Science and Psychology, University of Mohaghegh Ardabili, Ardabil, Iran

*Corresponding Author: AmirAli Jafarnezhadgero, Email: Amiralijafarnezhad@gmail.com

Abstract

Background: This research aimed to assess the effect of medicine ball exercise on the amplitude of electromyography activity in back pain patients while walking.

Methods: Twenty back pain patients participated in this research. The participants were divided into two intervention (age: 26.4 ± 2.00 years) and control (age: 28.8 ± 2.0 years) groups. The intervention involved training with medicine ball. The Roland Morris questionnaire was employed to measure the pain index. In addition, the participants walked at self-selected gait speed along an 18-m level walkway. During walking, muscle activities of the following muscles were recorded: gastrocnemius lateral (GAS-L), biceps femoris (BF), Semimembranosus muscle (SM), gluteus medius (GM), Erector spinae right (ES-R), Erector spinae left (ES-L), Internal abdominal oblique (IA-O) and External abdominal oblique (EA-O).

Results: Significant main effects of "Time" for disability index were observed ($P < 0.001$). Also, significantly lower disability index was observed after exercise with medicine ball compared with pre-intervention. Significant main effects of "Time" for BF ($P < 0.001$, $\eta^2 = 0.532$) and ES-R ($P < 0.001$, $\eta^2 = 0.449$) muscles activities during midstance phase were also found. The findings demonstrated lower BF activities after exercise with medicine ball compared to before it. Finally, greater ES-R activities post-exercise were observed with medicine ball compared with before it.

Conclusion: This study revealed that exercise with medicine ball improved disability index and muscle activities in individuals with back pain while walking.

Keywords: Walking, Electromyography, Pain

Citation: Fakhri Mirzang E, Jafarnezhadgero A, Bolboli L. Impact of medicine ball training on amplitude of electromyography activity in back pain patients during gait. *Journal of Kerman University of Medical Sciences*. 2025;32:2830. doi:10.34172/jkmu.2830

Received: April 9, 2021, **Accepted:** August 4, 2025, **ePublished:** August 10, 2025

Introduction

Low back pain (LBP) has a global prevalence of 6.8% (1), occurring in 19.6% of adults (2). People with LBP demonstrate earlier fatigue in trunk muscles (1, 3, 4). A study reported that patients with back pain do not have the ability to voluntarily recruit the muscular median frequency (5). Many studies have reported conflicting findings regarding muscular activity in LBP subjects (6-9). In a previous research, (10) used EMG to demonstrate that the lumbar paraspinal muscles are minimally active in relaxed standing.

Williams (1965) (11) recommended exercises to strengthen the abdominal muscles, while Cyriax (1976) (12) focused more on keeping the lumbar lordosis to manage persistent pain. Other physical methods and behavioral strategies should also be taken into account to properly treat the combination of symptoms, especially

when pain is a major factor (12). Researchers have researched the trunk muscles extensively using different methods to understand how these muscles malfunction. Surface EMG has been a suitable system for how trunk muscles work along certain movements and positions (3).

Meanwhile, rehabilitation is recommended as a key point for treatment of LBP (13). Sport activity that focus on strengthening the trunk and back muscles while improving body coordination may help individuals with chronic back pain recover more effectively (14, 15). In spite of massive research, the root causes of LBP. Resistance training can help improve key fitness markers such as muscle strength, explosive power, and endurance (16). The real advantage of these exercises is that they engage far more muscles and establish greater muscle activation compared to traditional strength exercises. Research by Wilk et al suggested use of unstable surfaces such as a medicine ball for similar



movements along advanced recovery stages to maximize benefits (17).

The author argues this is because these exercises force the body into movements that require far greater dynamic stability (18). For example, Norian et al demonstrated that rehabilitation with medicine ball improved knee co-contraction in LBP subjects during gait (19). Research on medicine ball exercises has examined how upper and lower body strength as well as power relate to overall explosive force during throws (20). However, there is still limited evidence backing their effectiveness. Existing studies are narrow in scope—often due to small sample sizes—and fail to measure critical factors such as muscle activity (EMG) in individuals with back pain, which could help predict injury risks.

There is no available study that has specifically investigated how medicine ball exercises affect individuals with lower back pain. Thus, this study aimed to ascertain the effect of medicine ball training on amplitude of electromyography activity in those with LBP during walking. We hypothesized that pain index in individuals with LBP would be smaller after exercise with medicine ball. We also hypothesized that muscle activities in individuals with LBP would diminish after exercise with medicine ball (21, 22).

Methods

Participants

Twenty LBP patients volunteered to participate in the present study. They were divided into two intervention (age: 26.4 ± 2.05 years, mass: 74.08 ± 3.06 kg, height: 181 ± 7.04 cm) and control (age: 28.8 ± 2.05 years, mass: 88.45 ± 3.01 kg, height: 181 ± 7.01 cm). The exclusion criteria included underlying pain of trauma, nerve or spinal cord injury in the lumbar spine, disc herniation, rheumatic disease, inflammation, previous lumbar surgery, pregnancy, as well as cardiorespiratory and metabolic diseases. Patients with LBP were included in the study after signing the consent form. The intervention group performed resistance exercises with medicine ball while the control group participated in pre-test and post-test only.

Pain and muscle activity assessment

The Roland Morris Questionnaire was employed to measure the quantity of pain (23). To record the reaction forces of the ground during walking before and after the training, the Bertec Force Plates (USA) with a sampling frequency of 1000 Hz was utilized to record kinetic data. A frequency cut-off of 20 Hz was applied to filter the reaction forces of the ground. The peak reaction forces and the time to reach the peak forces were extracted according to the study by Jafarnezhadgero et al (24).

An EMG system (Biometrics, UK, sample rate: 1000 Hz) with surface electrodes was employed to record EMG of the gastrocnemius lateral (GAS-L), biceps femoris

(BF), semimembranosus muscle (SM), gluteus medius (GM), erector spinae right (ES-R), erector spinae left (ES-L), internal abdominal oblique (IA-O), and external abdominal oblique (EA-O). Thereafter, the skin was abraded prior to electrode placement (25). For EMG analyses, the gait cycle included loading phase, the mid-stance, and push off phases (26). The maximum voluntary isometric contraction (MVIC) was recorded EMG data normalization (27).

Exercise program

The training was performed for three sessions each week on nonconsecutive days (30-32) (Table 1).

Statistical analyses

The normality of data was affirmed through the Shapiro-Wilk test. A mixed ANOVA with repeated measures was employed for analysis using SPSS 26.

Results

Anthropometric characteristics of two groups are reported in Table 2.

Significant time effects for SM activity during loading were observed ($P < .001$, $\eta^2 = 0.517$). Lower SM muscle activity was found after exercise with medicine ball compared with pre-exercise.

Significant time effects were observed for BF ($P < .001$, $\eta^2 = 0.532$) and ES-R ($P < .001$, $\eta^2 = 0.449$) muscle activity along the midstance phase. Lower BF activity was reported post-exercise with medicine ball. Also, greater ES-R muscle activity after exercise with medicine ball compared with pre-exercise was reported (Table 3).

Effects of "Group" for ES-L was significant ($P = 0.023$, $\eta^2 = 0.255$). Muscle activity during the midstance phase. Greater ES-L activity in the intervention than in the control group was reported (Table 3).

Group effects for ES-R ($P = 0.003$, $\eta^2 = 0.394$) and EA-O ($P = 0.006$, $\eta^2 = 0.350$) activities during the midstance phase was significant. Greater ES-R and EA-O activities in the intervention than in the control group was observed (Table 3).

Group-by-Time interactions were significant for IA-O muscle activity during loading phase ($P = 0.005$, $\eta^2 = 0.089$). Post hoc analysis demonstrated significantly greater IA-O muscle activity in the intervention group after exercise with medicine ball compared with pre-exercise (Table 3).

Group-by-Time interactions were significant for ES-R activities during midstance phase ($P = 0.012$, $\eta^2 = 0.300$). Post hoc analysis demonstrated greater ES-R activity in the intervention group after exercise with medicine ball compared pre-exercise (Table 3).

Discussion

This study was the first to explore the influence of medicine ball training on the amplitude of electromyography

Table 1. Exercise program with medicine ball

Week	Sections of each session	Intensity (Vo ₂ Max %)	Section time in each session (minutes)
Weeks 1: (three sessions per week)	Warm-up	50	10
	Specific isometric and isotonic exercises for the lower body and upper body	50	40
	Cooldown	50	10
Weeks 2: (three sessions per week)	Warm-up	50	10
	Specific isometric and isotonic exercises for the lower body and upper body	50	40
	Cooldown	50	10
Weeks 3: (three sessions per week)	Warm-up	55	10
	Specific isometric and isotonic exercises for the lower body and upper body	55	40
	Cooldown	55	10
Weeks 4: (three sessions per week)	Warm-up	60	10
	Specific isometric and isotonic exercises for the lower body and upper body	65-70	40
	Cooldown	60	10

Warm-up: Jogging, dynamic stretching movements

Main section: isotonic and isometric as well as cyclic training with beta medicine ball.

Cooldown: Local endurance and relaxing the muscles, static stretching

Table 2. Group-specific baseline values of all reported anthropometrics together (M±SD).

Parameter	Intervention group	Control group	Sig.
Age (years)	28.8±2.5	26.4±2.5	0.119
Body height (cm)	1.81±7.1	1.81±7.4	0.964
Body mass (kg)	88.45±3.1	74.08±3.6	0.208

Note. M, mean; SD, standard deviation; Sig, Significant

activity in back pain patients during gait.

The results demonstrated significantly lower SM muscle activity after exercise with medicine ball compared with pre-exercise. The results revealed significantly lower BF muscle activities post-exercise with medicine ball compared with pre-exercise. The hamstrings play a key role in stabilizing the knee joint and are necessary for maintaining dynamic control of the knee joint. Meanwhile, men tend to have higher levels of active and passive hamstring stiffness than women, which may help explain why knee-joint injuries are more frequently observed in female athletes (28). The attachment of the hamstring muscles to upper limbs suggested that active stiffness in this muscles might affect pelvic and core stability; nonetheless, the relationship is currently speculative and unsubstantiated (29).

The roles of hamstring muscles to these observed deteriorations are greater in individuals with LBP (30). When showed individuals with LBP is necessary to consider muscles and joints in the core muscles (31). Muscle groups surrounding the lower limb have been reported to have similar fatiguing patterns in individuals with LBP (32).

It is believed that LBP is associated during walking, increased trunk stiffness, higher activation of the erector spinae (33) and the hamstring muscles (29). The result of this study suggests a clinically important effect on hamstring function which occurs after exercise with medicine ball in patients with a history of LBP. This

change may be indicative of a delayed muscle response to the resistance training.

The finding demonstrated greater EA-O activities in the intervention group than in the control group. Exercises performed in different positions required similar levels of extensor muscle activation but differed in flexor muscle recruitment patterns, with the oblique muscles presenting greater engagement compared to the rectus abdominis. The oblique muscles and rectus abdominis work simultaneously to stabilize pelvic positioning, while the multifidus acts as the antagonist muscle along these movements (34).

Musculoskeletal pain leads to reduction of IA-O activity while the muscle as an agonist (35). Pain could result in compensatory actions from other muscles (36). Group-by-Time interactions for IA-O activities at loading were significant. Greater IA-O activities were observed in the intervention group after exercise with medicine ball compared with pre-exercise. These results are in accordance with previous studies confirming a reduction of activation in the right multifidus, iliocostalis, and abdominal muscles in LBP individuals during walking (37, 38). In comparison to other studies, where other resistance programs were used in individuals with LBP, it is difficult to generalize the conclusions from other resistance programs to the patients' population.

Several limitations should be noted. First, the small sample size limited our ability to detect smaller effects that might have reached statistical significance in a larger sample size. Further, the study exclusively included male participants, so we cannot generalize the results to the group of men and women. Also, in this study, the activity of some other muscles of the lower and upper limbs was not recorded owing to the limitations of installing the electrodes, which should be examined in future studies. Also, lack of registration of kinematic variables was another limitation of the present study.

Table 3. Muscle activity (% maximum voluntary isometric contraction [MVIC]) at the loading, midstance and push off phases along walking

Intervention group				Control group		SIG (Eta)		
Muscle activity		PRE	POST	PRE	POST	Time	group	Time*group
GAS LAT	L	12.14±9.56	11.60±5.20	12.64±13.34	15.85±17.38	0.746 (0.006)	0.526 (0.023)	0.649 (0.012)
	MS	16.32±17.33	24.3±12.60	19.17±18.44	34.74±24.40	0.116(0.132)	0.166(0.104)	0.290(0.062)
	PO	9.80±4.21	10.82±6.95	6.36±3.11	10.16±10.1	0.273(0.066)	0.342(0.050)	0.523(0.023)
BF	L	7.57±3.93	9.1±7.79	10.82±6.95	13.89±17.36	0.525(0.23)	0.198(0.090)	0.817(0.003)
	MS	30.65±19.18	10.41±7.52	45.19±32.37	10.40±7.26	<.001(0.532)	0.261(0.070)	0.247(0.074)
	PO	12.81±9.47	18.55±16.93	9.65±7.68	7.23±2.93	0.575(0.018)	0.067(0.175)	0.178(0.098)
SM	L	36.50±24.94	16.42±11.32	39.11±27.13	9.81±7.35	<.001(0.517)	0.776(0.005)	0.437(0.036)
	MS	20.61±10.56	18.20±13.70	18.19±13.29	15.89±16.69	0.636(0.013)	0.532(0.022)	0.991(0.000)
	PO	7.57±3.93	15.52±12.30	10.82±6.95	9.80±6.98	0.164(0.105)	0.658(0.011)	0.077(0.164)
GLUT MED	L	15.12±14.40	17.12±12.63	7.98±5.87	8.27±6.77	0.792(0.004)	0.055(0.200)	0.843(0.002)
	MS	15.61±13.05	15.50±13.85	13.07±9.38	9.44±7.5	0.584(0.017)	0.262(0.069)	0.607(0.015)
	PO	54.87±23.47	18.15±12.87	54.62±27.10	10.60±1.62	<.001(0.726)	0.291(0.062)	0.337(0.051)
ES-R	L	10.33±1.80	14.36±8.50	16.24±16.83	8.16±5.70	0.511(0.26)	0.968(0.000)	0.061(0.192)
	MS	7.60±2.74	25.36±14.63	6.34±4.55	9.18±5.63	0.001(0.449)	0.003(0.394)	0.012(0.300)
	PO	25.75±15.35	20.25±13.26	21.71±11.25	20.56±15.64	0.527(0.023)	0.618(0.014)	0.678(0.010)
ES-L	L	32.22±92.58	17.19±16.64	8.63±6.15	13.5±12.66	0.329(0.053)	0.023(0.255)	0.084(0.157)
	MS	26.80±22.67	14.35±12.99	21.62±15.56	13.95±7.80	0.096(0.146)	0.500(0.026)	0.682(0.010)
	PO	16.70±13.35	13.35±11.46	21.44±24.67	12.24±9.54	0.251(0.073)	0.707(0.008)	0.586(0.017)
IA-O	L	14.88±15.38	19.21±15.61	18.82.±17.96	19.80±13.84	0.642 (0.012)	0.702(0.008)	0.005(0.089)
	MS	17.72±11.1	22.8±12.38	15.28±13.15	11.84±6.58	0.875(0.001)	0.134(0.120)	0.190(0.093)
	PO	21.59±16.77	13.19±10.75	19.32±12.9	21.57±24.20	0.605(0.015)	0.526(0.023)	0.374(0.044)
EA-O	L	20.41±22.44	15.51±18.18	17.49±17.86	11.41±12.86	0.372(0.044)	0.529(0.022)	0.923(0.001)
	MS	34.69±23.71	17.31±11.42	13.15±5.61	15.12±10.52	0.157(0.108)	0.006(0.350)	0.080(0.160)
	PO	38.47±28.18	10.96±5.22	36.88±25.50	10.95±5.22	0.048(0.200)	0.967(0.000)	0.873(0.001)

Note. GAS LAT, Gastrocnemius lateral; BF, Biceps femoris; SM, Semimembranosus; GM, Gluteus medius; ES-R, Erector spinae right; ES-L, Erector spinae left; IA-O, Internal abdominal oblique; EA-O, External abdominal oblique
Bold number dimonstrit significant difference

Conclusion

This study revealed that exercise with medicine ball improve disability index and muscle activities in individuals with back pain during walking.

Acknowledgments

We gratefully acknowledge Mr. Ehsan Fakhri Mirzanag, Dr AmirAli Jafarnezhadgero, and Dr, Lotfali Bolboli for our help and all those who helped us in the implementation of this study

Authors’ Contribution:

Conceptualization: AmirAli Jafarnezhadgero.
Data curation: Ehsan Fakhri Mirzanag.
Formal analysis: Lotfali Bolboli.
Funding acquisition: AmirAli Jafarnezhadgero.
Investigation: AmirAli Jafarnezhadgero.
Methodology: Ehsan Fakhri Mirzanag.
Project administration: AmirAli Jafarnezhadgero.
Resources: Ehsan Fakhri Mirzanag.

Software: Ehsan Fakhri Mirzanag.
Supervision: Lotfali Bolboli.
Validation: Lotfali Bolboli.
Visualization: Lotfali Bolboli.
Writing–original draft: Ehsan Fakhri Mirzanag.
Writing–review & editing: Ehsan Fakhri Mirzanag.

Ethical Approval

The study was conducted in accordance with the latest version of the Declaration of Helsinki. Ethical approval was obtained from the local ethical committee (IR.ARUMS.REC.1397.031).

Funding

This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Gudavalli MR. Biomechanics research on flexion-distraction procedure. In: Cox JM, ed. Low Back Pain: Mechanisms,

- Diagnosis and Treatment. Philadelphia, PA: Lippincott Williams & Wilkins; 1999.
2. Meucci RD, Fassa AG, Faria NM. Prevalence of chronic low back pain: systematic review. *Rev Saude Publica*. 2015;49:1. doi: [10.1590/s0034-8910.2015049005874](https://doi.org/10.1590/s0034-8910.2015049005874).
3. De Luca CJ. Use of the surface EMG signal for performance evaluation of back muscles. *Muscle Nerve*. 1993;16(2):210-6. doi: [10.1002/mus.880160216](https://doi.org/10.1002/mus.880160216).
4. McGregor AH, Bull AM, Byng-Maddick R. A comparison of rowing technique at different stroke rates: a description of sequencing, force production and kinematics. *Int J Sports Med*. 2004;25(6):465-70. doi: [10.1055/s-2004-820936](https://doi.org/10.1055/s-2004-820936).
5. Greenough CG, Oliver CW, Jones AP. Assessment of spinal musculature using surface electromyographic spectral color mapping. *Spine (Phila Pa 1976)*. 1998;23(16):1768-74. doi: [10.1097/00007632-199808150-00009](https://doi.org/10.1097/00007632-199808150-00009).
6. Ahern DK, Follick MJ, Council JR, Laser-Wolston N, Litchman H. Comparison of lumbar paravertebral EMG patterns in chronic low back pain patients and non-patient controls. *Pain*. 1988;34(2):153-60. doi: [10.1016/0304-3959\(88\)90160-1](https://doi.org/10.1016/0304-3959(88)90160-1).
7. Arena JG, Sherman RA, Bruno GM, Young TR. Electromyographic recordings of low back pain subjects and non-pain controls in six different positions: effect of pain levels. *Pain*. 1991;45(1):23-8. doi: [10.1016/0304-3959\(91\)90160-y](https://doi.org/10.1016/0304-3959(91)90160-y).
8. Grabel JA. Electromyographic Study of Low Back Muscle Tension in Subjects with and Without Chronic Low Back Pain [dissertation]. ProQuest Information & Learning; 1973.
9. Kravitz E, Moore ME, Glaros A. Paralumbal muscle activity in chronic low back pain. *Arch Phys Med Rehabil*. 1981;62(4):172-6.
10. Floyd WF, Silver PH. The function of the erector spinae muscles in certain movements and postures in man. *J Physiol*. 1955;129(1):184-203. doi: [10.1113/jphysiol.1955.sp005347](https://doi.org/10.1113/jphysiol.1955.sp005347).
11. Williams PC. The Lumbosacral Spine: Emphasizing Conservative Management. McGraw-Hill; 1965.
12. Hendry NG. Textbook of orthopaedic medicine. Volume 1: Diagnosis of soft tissue lesions. James Cyriax, Sixth edition. 230×160 mm. Pp. 756+xii, with 204 illustrations. 1975. London: Baillière Tindall. £12.50. *Br J Surg*. 1976;63(8):671. doi: [10.1002/bjs.1800630826](https://doi.org/10.1002/bjs.1800630826).
13. Bekkering GE, Hendriks HJ, Koes BW, Oostendorp RA, Ostelo R, Thomassen JM, et al. Dutch physiotherapy guidelines for low back pain. *Physiotherapy*. 2003;89(2):82-96. doi: [10.1016/s0031-9406\(05\)60579-2](https://doi.org/10.1016/s0031-9406(05)60579-2).
14. George AJ, Kumar D, Nikhil NP. Effectiveness of trunk proprioceptive neuromuscular facilitation training in mechanical low back pain. *Int J Curr Res*. 2013;5(7):1965-8.
15. Kofotolis N, Kellis E. Effects of two 4-week proprioceptive neuromuscular facilitation programs on muscle endurance, flexibility, and functional performance in women with chronic low back pain. *Phys Ther*. 2006;86(7):1001-12. doi: [10.1093/ptj/86.7.1001](https://doi.org/10.1093/ptj/86.7.1001).
16. Petrofsky JS, Prowse M, Lemoine M, Bartelink T, Batt J, Suh HJ, et al. Muscle use during exercise with a mini medicine ball compared to other abdominal exercise modalities. *J Appl Res*. 2008;8(2):95-115.
17. Wilk R, Weizman I, Shilo BZ. Trachealess encodes a bHLH-PAS protein that is an inducer of tracheal cell fates in *Drosophila*. *Genes Dev*. 1996;10(1):93-102. doi: [10.1101/gad.10.1.93](https://doi.org/10.1101/gad.10.1.93).
18. de Oliveira AS, de Morais Carvalho M, de Brum DP. Activation of the shoulder and arm muscles during axial load exercises on a stable base of support and on a medicine ball. *J Electromyogr Kinesiol*. 2008;18(3):472-9. doi: [10.1016/j.jelekin.2006.09.009](https://doi.org/10.1016/j.jelekin.2006.09.009).
19. Norian E, Fakhri Mirzanag M, Jafarnejadgero A. Effect of exercise therapy with medicine ball on directed and general co-contraction of knee joint muscles in low back pain patients during walking. *J Anesth Pain*. 2023;13(4):1-11.
20. Ikeda Y, Kijima K, Kawabata K, Fuchimoto T, Ito A. Relationship between side medicine-ball throw performance and physical ability for male and female athletes. *Eur J Appl Physiol*. 2007;99(1):47-55. doi: [10.1007/s00421-006-0316-4](https://doi.org/10.1007/s00421-006-0316-4).
21. Alavi Mehr SM, Mousavi SH, Jafarnejadgero A. The effect of a selected exercise protocol on trunk and lower limb muscle activity of older adults with both low back pain and pronated feet during walking. *Sport Sci Health Res*. 2021;14(1):129-40. doi: [10.22059/sshr.2023.350097.1065](https://doi.org/10.22059/sshr.2023.350097.1065).
22. Pakzad M, Fung J, Preuss R. Pain catastrophizing and trunk muscle activation during walking in patients with chronic low back pain. *Gait Posture*. 2016;49:73-7. doi: [10.1016/j.gaitpost.2016.06.025](https://doi.org/10.1016/j.gaitpost.2016.06.025).
23. Mousavi SJ, Parnianpour M, Mehdian H, Montazeri A, Mobini B. The Oswestry Disability Index, the Roland-Morris Disability Questionnaire, and the Quebec Back Pain Disability Scale: translation and validation studies of the Iranian versions. *Spine (Phila Pa 1976)*. 2006;31(14):E454-9. doi: [10.1097/01.brs.0000222141.61424.f7](https://doi.org/10.1097/01.brs.0000222141.61424.f7).
24. Jafarnejadgero A, Fatollahi A, Amirzadeh N, Siahkhouhian M, Granacher U. Ground reaction forces and muscle activity while walking on sand versus stable ground in individuals with pronated feet compared with healthy controls. *PLoS One*. 2019;14(9):e0223219. doi: [10.1371/journal.pone.0223219](https://doi.org/10.1371/journal.pone.0223219).
25. Hermens HJ, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, et al. European recommendations for surface electromyography. Roessingh Research and Development. 1999;8(2):13-54.
26. Murley GS, Buldt AK, Trump PJ, Wickham JB. Tibialis posterior EMG activity during barefoot walking in people with neutral foot posture. *J Electromyogr Kinesiol*. 2009;19(2):e69-77. doi: [10.1016/j.jelekin.2007.10.002](https://doi.org/10.1016/j.jelekin.2007.10.002).
27. Farahpour N, Jafarnejadgero A, Allard P, Majlesi M. Muscle activity and kinetics of lower limbs during walking in pronated feet individuals with and without low back pain. *J Electromyogr Kinesiol*. 2018;39:35-41. doi: [10.1016/j.jelekin.2018.01.006](https://doi.org/10.1016/j.jelekin.2018.01.006).
28. Blackburn JT, Riemann BL, Padua DA, Guskiewicz KM. Sex comparison of extensibility, passive, and active stiffness of the knee flexors. *Clin Biomech (Bristol)*. 2004;19(1):36-43. doi: [10.1016/j.clinbiomech.2003.09.003](https://doi.org/10.1016/j.clinbiomech.2003.09.003).
29. Hanada EY, Johnson M, Hubley-Kozey C. A comparison of trunk muscle activation amplitudes during gait in older adults with and without chronic low back pain. *PM R*. 2011;3(10):920-8. doi: [10.1016/j.pmrj.2011.06.002](https://doi.org/10.1016/j.pmrj.2011.06.002).
30. Hart JM, Kerrigan DC, Fritz JM, Saliba EN, Gansneder B, Ingersoll CD. Contribution of hamstring fatigue to quadriceps inhibition following lumbar extension exercise. *J Sports Sci Med*. 2006;5(1):70-9.
31. Harris-Hayes M, Sahrmann SA, Van Dillen LR. Relationship between the hip and low back pain in athletes who participate in rotation-related sports. *J Sport Rehabil*. 2009;18(1):60-75. doi: [10.1123/jsr.18.1.60](https://doi.org/10.1123/jsr.18.1.60).
32. Kankaanpää M, Taimela S, Laaksonen D, Hänninen O, Airaksinen O. Back and hip extensor fatigability in chronic low back pain patients and controls. *Arch Phys Med Rehabil*. 1998;79(4):412-7. doi: [10.1016/s0003-9993\(98\)90142-3](https://doi.org/10.1016/s0003-9993(98)90142-3).
33. Vogt L, Pfeifer K, Banzer W. Neuromuscular control of walking with chronic low-back pain. *Man Ther*. 2003;8(1):21-8. doi: [10.1054/math.2002.0476](https://doi.org/10.1054/math.2002.0476).

34. Shields RK, Heiss DG. An electromyographic comparison of abdominal muscle synergies during curl and double straight leg lowering exercises with control of the pelvic position. *Spine (Phila Pa 1976)*. 1997;22(16):1873-9. doi: [10.1097/00007632-199708150-00012](https://doi.org/10.1097/00007632-199708150-00012).
35. Kenefati G, Rockholt MM, Ok D, McCartin M, Zhang Q, Sun G, et al. Changes in alpha, theta, and gamma oscillations in distinct cortical areas are associated with altered acute pain responses in chronic low back pain patients. *Front Neurosci*. 2023;17:1278183. doi: [10.3389/fnins.2023.1278183](https://doi.org/10.3389/fnins.2023.1278183).
36. Li W, Gong Y, Liu J, Guo Y, Tang H, Qin S, et al. Peripheral and central pathological mechanisms of chronic low back pain: a narrative review. *J Pain Res*. 2021;14:1483-94. doi: [10.2147/jpr.S306280](https://doi.org/10.2147/jpr.S306280).
37. Zhang Z, Zou J, Lu P, Hu J, Cai Y, Xiao C, et al. Analysis of lumbar spine loading during walking in patients with chronic low back pain and healthy controls: an OpenSim-Based study. *Front Bioeng Biotechnol*. 2024;12:1377767. doi: [10.3389/fbioe.2024.1377767](https://doi.org/10.3389/fbioe.2024.1377767).
38. Smith JA, Kulig K. Altered multifidus recruitment during walking in young asymptomatic individuals with a history of low back pain. *J Orthop Sports Phys Ther*. 2016;46(5):365-74. doi: [10.2519/jospt.2016.6230](https://doi.org/10.2519/jospt.2016.6230).