

Unusual Presentations of Cutaneous Leishmaniasis in an Endemic Area, Kerman Province, Iran

Simin Shamsi Meymandi, M.D.¹, Maryam Khalili, M.D.², Saman Mohammadi, M.D.²,
Mahin Aflatoonian, M.D.³, Rahim Ahmadi, M.D.⁴, Alireza Zeynadini Meymand, M.D.⁴

1- Professor of Dermatology, Pathology and Stem Cell Research Center, Kerman University of Medical Sciences, Kerman, Iran

2- Assistant Professor of Dermatology, Department of Dermatology, Afzalipour School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

3- Assistant Professor of Dermatology, Department of Dermatology, Afzalipour School of Medicine, Kerman University of Medical Sciences, Kerman, Iran (Corresponding author; maflatoonian@gmail.com)

4- Resident of Dermatology, Afzalipour School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Received: 1 July, 2017

Accepted: 3 November, 2017

ARTICLE INFO

Article type:

Case Report

Keywords:

Leishmaniasis
Sporotrichoid
Lichenoid
Eczematous

Abstract

Leishmaniasis is a infectious disease with broad spectrum of clinical features from typical noduloulcerative lesion to atypical lesions. In this article, we presented 4 cases with atypical presentation of cutaneous leishmaniasis lesions (including eczematous, sporotrichoid and lichenoid) in an endemic area in Iran, Kerman. The first case was a 28- year- old man presented with a pruritic purplish -red plaque with lichenoid appearance around proximal nail fold of right third finger since a few months ago. The second case was a young woman with an indurated yellowish-brown plaque with peripheral depigmentation firstly diagnosed as keloid with subsequent intralesional triamcinolone injection. The third case was a middle-aged woman with multiple ulcerated reddish-brown nodules and plaques in a linear array on both legs and one forearm (sporotrichoid pattern). The forth case was a man with a painful indurated reddish-brown plaque with yellowish crust and swelling resembling infected contact dermatitis.

Copyright: 2017 The Author(s); Published by Kerman University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Shamsi Meymandi S, Khalili M, Mohammadi S, Aflatoonian M, Ahmadi R, Zeynadini Meymand A.R. Unusual Presentations of Cutaneous Leishmaniasis in an Endemic Area, Kerman Province, Iran. Journal of Kerman University of Medical Sciences, 2017; 24(6): 527-531.

Introduction

Leishmaniasis is a common protozoan infectious disease which is transmitted by sand fly from phlebotomous family (1). Depending on parasite genetic background, number of parasite inoculation, host immune response to parasite the clinical presentation is classified to acute and chronic types (2).

Acute form may present with typical morphologies such as papule, plaque, ulcerative nodule and impetiginous (3). Atypical forms are classified based on their morphology and location. Atypical morphology can present as zosteriform, eczematous, erysipeloid, verrusiform and psoriasiform (4-7). Atypical locations include lips (macrocheilia), palmoplantar, periungual (paronychia) and genital area (chancriform) (8, 9).

Chronic type of leishmaniasis may present as lupoid form in site of previous scar from cutaneous leishmaniasis, or as a lesion with duration more than 1 year without any improvement (10).

In this article we describe 4 patients with uncommon presentation of cutaneous leishmaniasis.

Case reports

Case 1

A 28- year- old man presented with a history of a pruritic lesion around proximal nail fold of right third finger since a few months ago. Previously, the lesion was treated by general practitioner with impression of eczema with topical corticosteroid and emollient with no improvement. On physical examination we observed an indurated shiny purplish -red plaque with scale and minimal superficial ulcer. (Fig 1). Since, the patient was living in an endemic area of leishmaniasis treated with topical steroid without response, we decided to biopsy the lesion. Histologically, a granulomatous infiltration with numerous leishman bodies was reported. We referred our patient to leishmaniasis treatment center that treated with weekly Intralesional meglumine antimoniate in addition to cryotherapy every other week. The lesion was cleared after treatment course of 8 weeks.



Fig 1. An indurated shiny purplish -red plaque with scale and minimal ulcer on the surface

Case 2

An 18- year- old woman referred to dermatology center with a severely pruritic lesion on volar aspect of left forearm, since 1 year ago. Initially, she was treated by general practitioner with intralesional triamcinolone as keloid without any improvement. On examination, a 2 cm² indurated yellowish-brown plaque with apple-jelly appearance and some scales on the surface with depigmented margin was observed (Fig2). Due to the chronicity and induration of the lesion we performed skin biopsy, considering chronic or lupoid leishmaniasis, sarcoidosis and lupus vulgaris as differential diagnosis. After confirming the diagnosis of chronic leishmaniasis, the lesion was treated with combination of weekly intralesional meglumine antimoniate and biweekly cryotherapy for 3 months with complete cure.



Fig 2. An indurated yellowish-brown plaque with apple-jelly appearance and some scales on the surface and depigmentation at the margin of it

Case 3

A 53-year-old woman complaining of multiple ulcerated lesions in both legs and one forearm admitted in Afzalipour hospital in Kerman for further evaluation. In her past history she had suffered from diabetes mellitus taking insulin for 10 years. She had no other problem in her medical and drug history. On physical examination, no fever, lymphadenopathy or constitutional symptoms were detected. Multiple painless ulcerated reddish-brown nodules and plaques with crust formation in a linear array were observed (Figure 3).

Direct smear for bacterial infection was positive for pseudomonas which treated by ciprofloxacin. Skin biopsy from one of the lesions showed leishman bodies. The patient responded to intramuscular injection of meglumine antimoniate with dosage of 20 mg/kg/day for 3 weeks.



Figure 3. Multiple painless ulcerated reddish-brown nodules and plaques with crust formation on the surface in linear array

Case 4

A 47-year-old man with history of painful indurated plaque on dorsum of hand from 1 year ago referred to our clinic. Despite of treatment with combination of intralesional glucantime and cryotherapy for 4 weeks, the lesion continued to spread as indurated reddish-brown plaque with yellowish crust and swelling resembling infected contact dermatitis (Fig 4). Bacterial smear and culture from the lesion was positive and patient received intravenous cefazoline 1g, twice a day. After remission of bacterial infection, as induration of the lesion persisted, skin biopsy was performed that confirmed leishmaniasis. So he was treated with parenteral administration of glucantime for 3 weeks and oral allopurinol for 3 months.



Fig4. An indurated reddish-brown plaque with yellowish crust and multiple ulcers on the surface and swelling of hand resemble to infected contact dermatitis

Discussion

Atypical forms of cutaneous leishmaniasis can be seen with alteration (both increased and decreased) in host immune response. Cutaneous leishmaniasis in patients with cell mediated immunodeficiency secondary to HIV infection, diabetes mellitus, organ transplantation and immunosuppressive drugs can present as diffuse, disseminated, lupoid and sporotrichoid forms (10-12).

Sporotrichoid pattern is one of the uncommon presentations of CL characterized by spreading of the infection along the lymphatic drainage system. Other causes of the sporotrichoid lymphocutaneous infection are sporotrichosis schenckii, mycobacterium marinum, nocardia brasiliensis, bacillus anthracis, staphylococcus aureus and histoplasma capsulatum (13, 14).

Trauma due to intralesional injections, and immune system failure are the most common hypotheses proposed for sporotrichoid pattern of the infection (15).

Currently, first choice treatment of sporotrichoid pattern of cutaneous leishmaniasis is systemic therapy with meglumine

antimoniate. In case of contraindication or intolerance, antifungal drugs such as itraconazole can be used (15, 16).

Two out of 4 of our cases (case 2 and 4) have a history of treatment with local (topical and intralesional) corticosteroid. There are some reports of reactivation and/or alteration in clinical features of cutaneous leishmaniasis after topical and systemic corticosteroid that can be explained by local immunosuppression and anti-inflammatory effects (17).

Eczematous type is one of the atypical presentations of CL reported in previous articles. Occasionally, lesions are clinically similar to eczema and only biopsy and parasitological studies may lead to correct diagnosis. Although the exact mechanism of this form of CL isn't completely revealed, exaggerated cell-mediated immune response secondary to parasite invasion into the skin has been proposed (18, 19).

To date, there are several reports of leishmaniasis involving periungual area (20, 21), but our case (case 4) had a shiny and purplish lichenoid appearance with moderate pruritus. This can be due to repeated contact with irritant and allergic substance. According to our knowledge, this is the first report of lichenoid form of the leishmaniasis mimicking lichen simplex chronicus.

Conclusion

Atypical forms of cutaneous leishmaniasis can be seen in patients with cell mediated immunodeficiency or after application of topical and systemic corticosteroid. So we must consider atypical cutaneous leishmaniasis in differential diagnosis of indurated skin lesions unresponsive to usual treatments in endemic regions.

References

1. Bailey MS, Lockwood DN. Cutaneous leishmaniasis. *Clin Dermatol* 2007; 25(2):203-11.
2. Grevelink SA, Lerner EA. Leishmaniasis. *Journal of the American Academy of Dermatology* 1996; 34(2 Pt 1):257-72.
3. Wolff K, Johnson RA, Saavedra D. *Color Atlas & Synopsis of Clinical Dermatology*. 7th ed. New York: McGraw-Hill; 2013.
4. Bienzle U, Ebert F, Dietrich M. Cutaneous leishmaniasis in Eastern Saudi Arabia. Epidemiological and clinical features in a nonimmune population living in an endemic area. *Tropenmed Parasitol* 1978; 29(2):188-93.
5. Momeni A, Aminjavaheri M. Clinical picture of cutaneous leishmaniasis in Isfahan, Iran. *International Journal of Dermatology* 1994; 33(4):260-5.
6. Karıncaoglu Y, Esrefoglu M, Ozcan H. Atypical clinical form of cutaneous leishmaniasis: erysipeloid form. *International Journal of Dermatology* 2004; 43(11): 827-9.
7. Rubio F, Robayna G, Heranz P, Torres E, Peña J, Contreras F, et al. Leishmaniasis presenting as a psoriasiform eruption in AIDS. *British Journal of Dermatology* 1997; 136(5): 792-4.
8. El-Hoshy K. Lip leishmaniasis. *Journal of the American Academy of Dermatology* 1993; 28(4):661-2.
9. Raja KM, Khan AA, Hameed A, Rahman SB. Unusual clinical variants of cutaneous leishmaniasis in Pakistan. *Br J Dermatol* 1998; 139(1):111-3.
10. Akilov OE, Khachemoune A, Hasan T. Clinical manifestations and classification of Old World cutaneous leishmaniasis. *Int J Dermatol* 2007; 46(2):132-42.
11. Moravvej H, Barzegar M, Nasiri S, Abolhasani E, Mohebbi M. Cutaneous leishmaniasis with unusual clinical and histological presentation: report of four cases. *Acta Medica Iranica* 2013; 51(4):274-8.
12. Ghorbani GH, Hosseini SJ, Pourfarziani V, Ameli J, Akbari MH. Cutaneous leishmaniasis before and after renal transplantation; a case report. *Iranian Journal of Clinical Infectious Diseases* 2007; 2(1):33-6.
13. Ayatollahi J. Sporotrichoid cutaneous leishmaniasis in central Iran. *Iran J Med Sci* 2006; 31(3): 173-5.
14. Tobin EH, Jih WW. Sporotrichoid lymphocutaneous infections: etiology, diagnosis and therapy. *Am Fam Physician* 2001; 63(2):326-32.
15. Cozzani E, Satta R, Fausti V, Cottoni F, Parodi A. Cutaneous sporotrichoid leishmaniasis resistant to pentavalent antimonial therapy: complete resolution with itraconazole. *Clin Exp Dermatol* 2011; 36(1):49-51.
16. Morelli P, Gianelli E, Calattini S, Corbellino M, Antinori S, Meroni L. Itraconazole can be effective in the treatment of sporotrichoid leishmaniasis. *Journal of Travel Medicine* 2004; 11(5):328-30.
17. Marovich MA, Lira R, Shepard M, Fuchs GH, Kruezer R, Nutman TB, et al. Leishmaniasis recidivans recurrence after 43 years: a clinical and immunologic report after successful treatment. *Clin Infect Dis* 2001; 33(7):1076-9.
18. Nasiri S, Robati RM, Marefat A, Saeedi M, Sarrafi-rad N. Bilateral cutaneous leishmaniasis presenting as eczema-like eruptions on the hands. *Indian J Dermatol Venereol Leprol* 2010; 76(3):307.
19. Ayatollahi J, Bafghi AF, Shahcheraghi SH. Rare variants of cutaneous leishmaniasis presenting as eczematous lesions. *Med J Islam Repub Iran* 2014; 28: 71.
20. Chaabane H, Turki H. Cutaneous Leishmaniasis with a Paronychia-like Lesion. *N Engl J Med* 2014; 371:1736.
21. Mahmood K. An unusual paronychia. *Clinical and Experimental Dermatology* 2007; 32(5):611-2.